

Mahler measures and multiple Eisenstein values

François Brunault
UMPA – ÉNS Lyon

23 February 2024

Definition (Mahler, 1962)

For $P \in \mathbf{C}[x_1, \dots, x_n] \setminus \{0\}$, define

$$m(P) = \frac{1}{(2\pi i)^n} \int_{T^n} \log |P(x_1, \dots, x_n)| \frac{dx_1}{x_1} \cdots \frac{dx_n}{x_n}$$

where $T^n: |x_1| = \dots = |x_n| = 1$ is the n -torus.

Definition (Mahler, 1962)

For $P \in \mathbf{C}[x_1, \dots, x_n] \setminus \{0\}$, define

$$m(P) = \frac{1}{(2\pi i)^n} \int_{T^n} \log |P(x_1, \dots, x_n)| \frac{dx_1}{x_1} \cdots \frac{dx_n}{x_n}$$

where $T^n: |x_1| = \dots = |x_n| = 1$ is the n -torus.

- ▶ The integral converges absolutely.
- ▶ If P has coefficients in $\overline{\mathbf{Q}}$ then $m(P)$ should be a period in the sense of Kontsevich–Zagier.
- ▶ $m(P)$ measures the “size” of a polynomial in $\mathbf{Z}[x_1, \dots, x_n]$.
- ▶ Lehmer's problem (1933): For $P \in \mathbf{Z}[x]$ monic irreducible, not cyclotomic, can $m(P) > 0$ be arbitrarily small?

Theorem (Jensen, 1899)

For $P \in \mathbf{C}[x] \setminus \{0\}$, $P = a_d \prod_{i=1}^d (x - \alpha_i)$, we have

$$m(P) = \log |a_d| + \sum_{\substack{k=1 \\ |\alpha_k| \geq 1}}^d \log |\alpha_k|.$$

Theorem (Jensen, 1899)

For $P \in \mathbf{C}[x] \setminus \{0\}$, $P = a_d \prod_{i=1}^d (x - \alpha_i)$, we have

$$m(P) = \log |a_d| + \sum_{\substack{k=1 \\ |\alpha_k| \geq 1}}^d \log |\alpha_k|.$$

- ▶ Jensen's formula is still useful for multivariate polynomials: it reduces an n -dimension integral to an $(n-1)$ -dimensional one.
- ▶ Example: using Jensen's formula with respect to y , we have

$$m(1+x+y) = \frac{1}{2\pi i} \int_{\substack{|x|=1 \\ |1+x| \geq 1}} \log |1+x| \frac{dx}{x} = \frac{1}{2\pi} \int_{-2\pi/3}^{2\pi/3} \log |1+e^{i\theta}| d\theta$$

- ▶ How to evaluate further?

Timeline of identities

Smyth (1981): $m(1 + x + y) = \frac{3\sqrt{3}}{4\pi} L(\chi_3, 2)$

Here $L(\chi_3, s) = \sum_{n=1}^{\infty} \chi_3(n)/n^s$ is the Dirichlet L -function for

$$\chi_3(n) = \begin{cases} 1 & \text{if } n \equiv 1 \pmod{3} \\ -1 & \text{if } n \equiv 2 \pmod{3} \\ 0 & \text{if } n \equiv 0 \pmod{3} \end{cases}$$

Timeline of identities

Smyth (1981): $m(1 + x + y) = \frac{3\sqrt{3}}{4\pi} L(\chi_3, 2)$

Here $L(\chi_3, s) = \sum_{n=1}^{\infty} \chi_3(n)/n^s$ is the Dirichlet L -function for

$$\chi_3(n) = \begin{cases} 1 & \text{if } n \equiv 1 \pmod{3} \\ -1 & \text{if } n \equiv 2 \pmod{3} \\ 0 & \text{if } n \equiv 0 \pmod{3} \end{cases}$$

The proof uses the series expansion

$$\log |1 + e^{i\theta}| = -\operatorname{Re} \sum_{n=1}^{\infty} \frac{e^{-in\theta}}{n}.$$

and then integration from $\theta = -2\pi/3$ to $2\pi/3$.

Timeline of identities

Smyth (1981): $m(1 + x + y + z) = \frac{7}{2\pi^2} \zeta(3)$

Boyd and Deninger (1997):

$$m\left(x + \frac{1}{x} + y + \frac{1}{y} + 1\right) \stackrel{?}{=} \frac{15}{4\pi^2} L(E, 2) = L'(E, 0)$$

where $L(E, s) = \sum_{n=1}^{\infty} \frac{a_n}{n^s}$ is the L -function of the elliptic curve

$$E : x + \frac{1}{x} + y + \frac{1}{y} + 1 = 0.$$

Timeline of identities

Smyth (1981): $m(1 + x + y + z) = \frac{7}{2\pi^2} \zeta(3)$

Boyd and Deninger (1997):

$$m\left(x + \frac{1}{x} + y + \frac{1}{y} + 1\right) \stackrel{?}{=} \frac{15}{4\pi^2} L(E, 2) = L'(E, 0)$$

where $L(E, s) = \sum_{n=1}^{\infty} \frac{a_n}{n^s}$ is the L -function of the elliptic curve

$$E : x + \frac{1}{x} + y + \frac{1}{y} + 1 = 0.$$

- ▶ Discovered using numerical experiments + theoretical insights.
- ▶ Proved by Rogers and Zudilin (2011).

Boyd (1998): Families of conjectural identities, such as

$$m\left(x + \frac{1}{x} + y + \frac{1}{y} + k\right) \stackrel{?}{=} c_k L'(E_k, 0) \quad (k \in \mathbf{Z}, k \neq 0, \pm 4)$$

for some rational number $c_k \in \mathbf{Q}^\times$.

Boyd (1998): Families of conjectural identities, such as

$$m\left(x + \frac{1}{x} + y + \frac{1}{y} + k\right) \stackrel{?}{=} c_k L'(E_k, 0) \quad (k \in \mathbf{Z}, k \neq 0, \pm 4)$$

for some rational number $c_k \in \mathbf{Q}^\times$.

- ▶ Generalises to other families $m(P(x, y) + k)$ where the Newton polygon of $P(x, y)$ has $(0, 0)$ as the only interior point.
- ▶ Only finitely many such identities are proved.
- ▶ Related to the algebraic K -group $K_2(E_k)$ and the Bloch-Beilinson regulator map $K_2(E_k) \rightarrow \mathbf{R}$.

Conjecture (Boyd and Rodriguez Villegas, 2003):

$$m((1+x)(1+y)+z) \stackrel{?}{=} \frac{15^2}{4\pi^4} L(E, 3) = -2L'(E, -1)$$

where E is an elliptic curve of conductor 15.

Conjecture (Boyd and Rodriguez Villegas, 2003):

$$m((1+x)(1+y)+z) \stackrel{?}{=} \frac{15^2}{4\pi^4} L(E, 3) = -2L'(E, -1)$$

where E is an elliptic curve of conductor 15.

- ▶ There are several other $L(E, 3)$ identities, but they do not seem to come in families.
- ▶ Why does an elliptic curve appear here?

Conjecture (Boyd and Rodriguez Villegas, 2003):


$$m((1+x)(1+y)+z) \stackrel{?}{=} \frac{15^2}{4\pi^4} L(E, 3) = -2L'(E, -1)$$

where E is an elliptic curve of conductor 15.

- ▶ There are several other $L(E, 3)$ identities, but they do not seem to come in families.
- ▶ Why does an elliptic curve appear here?
- ▶ Because

$$E : \begin{cases} (1+x)(1+y)+z=0 \\ (1+\frac{1}{x})(1+\frac{1}{y})+\frac{1}{z}=0. \end{cases}$$

- ▶ Note that $\{(1+x)(1+y)+z=0\} \cap T^3 \subset E$.



Theorem (B. 2023)

We have $m((1+x)(1+y)+z) = -2L'(E, -1)$.

Theorem (B. 2023)

We have $m((1+x)(1+y)+z) = -2L'(E, -1)$.

- ▶ Now related to the K -group $K_4(E)$.
- ▶ Uses joint work with Zudilin on K_4 regulators.
- ▶ Key tool: Multiple modular values

$$\int_0^\infty f_1(iy_1)y_1^{s_1-1} dy_1 \int_{y_1}^\infty f_2(iy_2)y_2^{s_2-1} \dots \int_{y_{n-1}}^\infty f_n(iy_n)y_n^{s_n-1} dy_n$$

where f_1, \dots, f_n are modular forms.

Let $P = (1+x)(1+y) + z$.

Step 1: Deninger's method

Use Jensen's formula with respect to z .

$$\rightsquigarrow m(P) = \frac{1}{(2\pi i)^2} \int_{\Gamma} \eta(x, y, z)$$

where:

- ▶ η is an explicit closed 2-form on $V_P = \{P(x, y, z) = 0\}$.
- ▶ $\Gamma = \{(x, y, z) \in V_P : |x| = |y| = 1, |z| \geq 1\}$.

Step 2: Stokes's theorem

In our case, the form η happens to be *exact*. Writing $\eta = d\rho$, we have by Stokes's theorem

$$m(P) = \frac{1}{(2\pi i)^2} \int_{\Gamma} d\rho = \frac{1}{(2\pi i)^2} \int_{\gamma} \rho$$

with

$$\gamma = \partial\Gamma = \{(x, y, z) \in V_P : |x| = |y| = |z| = 1\}.$$

- ▶ $\gamma = V_P \cap T^3$ is contained in E .
- ▶ ρ is a *closed* 1-form on E .
- ▶ So we now have a 1-dimensional integral on E .

For any two functions f, g on E , define

$$\rho(f, g) = -D(f) \operatorname{darg}(g) + \frac{1}{3} \log |g| (\log |1-f| \operatorname{dlog} |f| - \log |f| \operatorname{dlog} |1-f|)$$

where $D: \mathbf{P}^1(\mathbf{C}) \rightarrow \mathbf{R}$ is the Bloch-Wigner dilogarithm

$$D(z) = \operatorname{Im} \left(\sum_{n=1}^{\infty} \frac{z^n}{n^2} \right) + \log |z| \operatorname{arg}(1-z).$$

Theorem (Lalín, 2015)

- ▶ $\rho = \rho(-y, x) - \rho(-x, y)$.
- ▶ γ is a generator of $H_1(E(\mathbf{C}), \mathbf{Z})^+$.

Step 3: Translate in the modular world

The elliptic curve E is isomorphic to the modular curve $X_1(15)$.

$$X_1(N) = \Gamma_1(N) \backslash \mathcal{H} \cup \{\text{cusps}\}$$

where

$$\Gamma_1(N) = \left\{ \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in \mathrm{SL}_2(\mathbf{Z}) : a, d \equiv 1 \pmod{N}, c \equiv 0 \pmod{N} \right\}.$$

Nice feature: the functions x and y on E correspond to *modular units* on $X_1(15)$, that is, all their zeros and poles are at the cusps.

Key fact: if u is a modular unit, then $d\log(u) = E_2(z)dz$ where E_2 is an Eisenstein series of weight 2.

We want to understand

$$\rho(u, v) = -D(u) \operatorname{darg}(v) + \frac{1}{3} \log |v| (\log |1-u| \operatorname{dlog} |u| - \log |u| \operatorname{dlog} |1-u|).$$

when u and v are modular units on $X_1(N)$.

We want to understand

$$\rho(u, v) = -D(u) \operatorname{darg}(v) + \frac{1}{3} \log |v| (\log |1-u| \operatorname{dlog} |u| - \log |u| \operatorname{dlog} |1-u|).$$

when u and v are modular units on $X_1(N)$.

- ▶ $\operatorname{dlog}(u)$ and $\operatorname{dlog}(v)$ are Eisenstein series, so the log terms of the formula are well-understood.

We want to understand

$$\rho(u, v) = -D(u) \operatorname{darg}(v) + \frac{1}{3} \log |v| (\log |1-u| \operatorname{dlog} |u| - \log |u| \operatorname{dlog} |1-u|).$$

when u and v are modular units on $X_1(N)$.

- ▶ $\operatorname{dlog}(u)$ and $\operatorname{dlog}(v)$ are Eisenstein series, so the log terms of the formula are well-understood.
- ▶ The challenging piece is $D(u)$. We use

$$d(D(u)) = \log |u| \operatorname{darg}(1-u) - \log |1-u| \operatorname{darg}(u)$$

- ▶ If u and $1-u$ are modular units, then $D(u)$ is an *iterated integral* of Eisenstein series.

Definition

For $k \geq 1$ and $\mathbf{x} = (x_1, x_2) \in (\mathbf{Z}/N\mathbf{Z})^2$, define the Eisenstein series

$$E_{\mathbf{x}}^{(k)}(\tau) = \sum_{m, n \in \mathbf{Z}} \frac{\exp\left(\frac{2\pi i}{N}(mx_2 - nx_1)\right)}{(m\tau + n)^k} \in M_k(\Gamma(N))$$

For $\mathbf{x}, \mathbf{y}, \mathbf{z} \in (\mathbf{Z}/N\mathbf{Z})^2$, define the *multiple Eisenstein values* (Manin, Brown)

$$\Lambda(\mathbf{x}, \mathbf{y}) := \int_0^{i\infty} E_{\mathbf{x}}^{(2)}(\tau_1) d\tau_1 \int_{\tau_1}^{i\infty} E_{\mathbf{y}}^{(2)}(\tau_2) d\tau_2$$

$$\Lambda(\mathbf{x}, \mathbf{y}, \mathbf{z}) := \int_0^{i\infty} E_{\mathbf{x}}^{(2)}(\tau_1) d\tau_1 \int_{\tau_1}^{i\infty} E_{\mathbf{y}}^{(2)}(\tau_2) d\tau_2 \int_{\tau_2}^{i\infty} E_{\mathbf{z}}^{(2)}(\tau_3) d\tau_3.$$

↪ The Mahler measure of P can be written as an explicit linear combination of multiple Eisenstein values.

Theorem (B.–Zudilin, 2023)

Let $\mathbf{x}, \mathbf{y}, \mathbf{z} \in (\mathbf{Z}/N\mathbf{Z})^2$ such that $\mathbf{x} + \mathbf{y} + \mathbf{z} = 0$. If all the coordinates of $\mathbf{x}, \mathbf{y}, \mathbf{z}$ are non-zero, then

$$\begin{aligned} & \operatorname{Re}(\Lambda(\mathbf{x}, \mathbf{y}, \mathbf{y}) - \Lambda(\mathbf{z}, \mathbf{y}, \mathbf{y}) + \Lambda(\mathbf{y}, \mathbf{x}, \mathbf{x}) - \Lambda(\mathbf{z}, \mathbf{x}, \mathbf{x}) + \Lambda(\mathbf{z}, \mathbf{y}, \mathbf{x}) + \Lambda(\mathbf{z}, \mathbf{x}, \mathbf{y}) \\ & \quad - (\Lambda(\mathbf{y}) - \Lambda(\mathbf{x}))(\Lambda(\mathbf{x}, \mathbf{y}) + \Lambda(\mathbf{y}, \mathbf{z}) + \Lambda(\mathbf{z}, \mathbf{x}))) = L'(F_{\mathbf{x}, \mathbf{y}}, -1) + c_{\mathbf{x}, \mathbf{y}} \zeta(3) \end{aligned}$$

for some explicit $F_{\mathbf{x}, \mathbf{y}} \in M_2(\Gamma(N))$, and $c_{\mathbf{x}, \mathbf{y}} \in \mathbf{Q}$.

Theorem (B.–Zudilin, 2023)

Let $\mathbf{x}, \mathbf{y}, \mathbf{z} \in (\mathbf{Z}/N\mathbf{Z})^2$ such that $\mathbf{x} + \mathbf{y} + \mathbf{z} = 0$. If all the coordinates of $\mathbf{x}, \mathbf{y}, \mathbf{z}$ are non-zero, then

$$\begin{aligned} & \operatorname{Re}(\Lambda(\mathbf{x}, \mathbf{y}, \mathbf{y}) - \Lambda(\mathbf{z}, \mathbf{y}, \mathbf{y}) + \Lambda(\mathbf{y}, \mathbf{x}, \mathbf{x}) - \Lambda(\mathbf{z}, \mathbf{x}, \mathbf{x}) + \Lambda(\mathbf{z}, \mathbf{y}, \mathbf{x}) + \Lambda(\mathbf{z}, \mathbf{x}, \mathbf{y}) \\ & \quad - (\Lambda(\mathbf{y}) - \Lambda(\mathbf{x}))(\Lambda(\mathbf{x}, \mathbf{y}) + \Lambda(\mathbf{y}, \mathbf{z}) + \Lambda(\mathbf{z}, \mathbf{x}))) = L'(F_{\mathbf{x}, \mathbf{y}}, -1) + c_{\mathbf{x}, \mathbf{y}} \zeta(3) \end{aligned}$$

for some explicit $F_{\mathbf{x}, \mathbf{y}} \in M_2(\Gamma(N))$, and $c_{\mathbf{x}, \mathbf{y}} \in \mathbf{Q}$.

Proving this formula requires two ingredients:

- ▶ **Interpolate** the multiple Eisenstein values to continuous parameters, viewing $(\mathbf{Z}/N\mathbf{Z})^2$ inside $(\mathbf{R}/\mathbf{Z})^2$ using $(x_1, x_2) \mapsto (\frac{x_1}{N}, \frac{x_2}{N})$.
- ▶ **Differentiate** with respect to these parameters to reduce the length of the iterated integrals.

Key lemma

For $\mathbf{x} = (x_1, x_2) \in (\mathbf{R}/\mathbf{Z})^2$, we have

$$\frac{d}{dx_2} E_{\mathbf{x}}^{(2)}(\tau) = \frac{d}{d\tau} E_{\mathbf{x}}^{(1)}(\tau).$$

Key lemma

For $\mathbf{x} = (x_1, x_2) \in (\mathbf{R}/\mathbf{Z})^2$, we have

$$\frac{d}{dx_2} E_{\mathbf{x}}^{(2)}(\tau) = \frac{d}{d\tau} E_{\mathbf{x}}^{(1)}(\tau).$$

So for example

$$\begin{aligned} \frac{d}{dy_2} \Lambda(\mathbf{x}, \mathbf{y}) &= \int_0^{i\infty} E_{\mathbf{x}}^{(2)}(\tau_1) d\tau_1 \int_{\tau_1}^{i\infty} \frac{d}{dy_2} E_{\mathbf{y}}^{(2)}(\tau_2) d\tau_2 \\ &= \int_0^{i\infty} E_{\mathbf{x}}^{(2)}(\tau_1) d\tau_1 \int_{\tau_1}^{i\infty} \frac{d}{d\tau_2} E_{\mathbf{y}}^{(1)}(\tau_2) d\tau_2 \\ &= \int_0^{i\infty} E_{\mathbf{x}}^{(2)}(\tau_1) (E_{\mathbf{y}}^{(1)}(i\infty) - E_{\mathbf{y}}^{(1)}(\tau_1)) d\tau_1. \end{aligned}$$

This reduces a double integral to a single integral.

To prove the formula

$$\begin{aligned} & \operatorname{Re}(\Lambda(x, y, y) - \Lambda(z, y, y) + \Lambda(y, x, x) - \Lambda(z, x, x) + \Lambda(z, y, x) + \Lambda(z, x, y) \\ & - (\Lambda(y) - \Lambda(x))(\Lambda(x, y) + \Lambda(y, z) + \Lambda(z, x))) = L'(F_{x,y}, -1) + c_{x,y}\zeta(3) \end{aligned}$$

we differentiate the LHS with respect to x_2 .

We get a sum of double integrals of the form

$$\int_0^{i\infty} E_{\mathbf{a}}^{(2)}(\tau_1) d\tau_1 \int_{\tau_1}^{i\infty} E_{\mathbf{b}}^{(2)}(\tau_2) E_{\mathbf{c}}^{(1)}(\tau_2) d\tau_2.$$

To prove the formula

$$\begin{aligned} & \operatorname{Re}(\Lambda(x, y, y) - \Lambda(z, y, y) + \Lambda(y, x, x) - \Lambda(z, x, x) + \Lambda(z, y, x) + \Lambda(z, x, y) \\ & \quad - (\Lambda(y) - \Lambda(x))(\Lambda(x, y) + \Lambda(y, z) + \Lambda(z, x))) = L'(F_{x,y}, -1) + c_{x,y}\zeta(3) \end{aligned}$$

we differentiate the LHS with respect to x_2 .

We get a sum of double integrals of the form

$$\int_0^{i\infty} E_{\mathbf{a}}^{(2)}(\tau_1) d\tau_1 \int_{\tau_1}^{i\infty} E_{\mathbf{b}}^{(2)}(\tau_2) E_{\mathbf{c}}^{(1)}(\tau_2) d\tau_2.$$

Miracle: The (complicated) linear combination of products

$E_{\mathbf{b}}^{(2)} E_{\mathbf{c}}^{(1)}$ is actually an Eisenstein series of weight 3!

This means that we have a *double Eisenstein value*.

The double Eisenstein values can be computed using the *Rogers-Zudilin method*. We get

$$\frac{d}{dx_2}(\text{LHS}) = \text{sum of } L\text{-values } L'(G_{\mathbf{a}}^{(1)} G_{\mathbf{b}}^{(2)}, 0)$$

for some (other) Eisenstein series $G^{(1)}$ and $G^{(2)}$.

The double Eisenstein values can be computed using the *Rogers-Zudilin method*. We get

$$\frac{d}{dx_2}(\text{LHS}) = \text{sum of } L\text{-values } L'(G_{\mathbf{a}}^{(1)} G_{\mathbf{b}}^{(2)}, 0)$$

for some (other) Eisenstein series $G^{(1)}$ and $G^{(2)}$.

This can be integrated to

$$\text{LHS} = \text{sum of } L\text{-values } L'(G_{\mathbf{a}}^{(1)} G_{\mathbf{b}}^{(1)}, -1)$$

We arrive at our L -value $L'(F_{\mathbf{x}, \mathbf{y}}, -1)$.

The double Eisenstein values can be computed using the *Rogers-Zudilin method*. We get

$$\frac{d}{dx_2}(\text{LHS}) = \text{sum of } L\text{-values } L'(G_{\mathbf{a}}^{(1)} G_{\mathbf{b}}^{(2)}, 0)$$

for some (other) Eisenstein series $G^{(1)}$ and $G^{(2)}$.

This can be integrated to

$$\text{LHS} = \text{sum of } L\text{-values } L'(G_{\mathbf{a}}^{(1)} G_{\mathbf{b}}^{(1)}, -1)$$

We arrive at our L -value $L'(F_{\mathbf{x}, \mathbf{y}}, -1)$.

Remark

We have no good understanding of the $\zeta(3)$ term in the formula.

The proof of the theorem also builds on:

- ▶ The Siegel modular units g_x for $x \in (\mathbf{Z}/N\mathbf{Z})^2$ on the modular curve $Y(N) = \Gamma(N)\backslash\mathcal{H}$
- ▶ Milnor symbols $\{g_x, g_y\}$ in $K_2(Y(N)) \otimes \mathbf{Q}$
- ▶ Three-term relations: if $x + y + z = 0$ then

$$\{g_x, g_y\} + \{g_y, g_z\} + \{g_z, g_x\} = 0.$$

- ▶ We can actually find a “triangulation”

$$g_x \wedge g_y + g_y \wedge g_z + g_z \wedge g_x = \sum_i m_i \cdot u_i \wedge (1 - u_i)$$

where u_i and $1 - u_i$ are modular units, and $m_i \in \mathbf{Q}$.

- ▶ This triangulation leads to an element of $K_4(Y(N)) \otimes \mathbf{Q}$.

This should extend in higher weight: for $k \geq 0$ and $\mathbf{x} \in (\mathbf{Z}/N\mathbf{Z})^2$, there is the *Eisenstein symbol*

$$\text{Eis}^k(\mathbf{x}) \in K_{k+1}(E(N)^k) \otimes \mathbf{Q}$$

where $E(N)^k$ is the k -fold fibre product of the universal elliptic curve $E(N)$ over the modular curve $Y(N)$.

Definition

For $k, \ell \geq 0$ and $\mathbf{x}, \mathbf{y} \in (\mathbf{Z}/N\mathbf{Z})^2$, define

$$X^k Y^\ell(\mathbf{x}, \mathbf{y}) = p_1^* \text{Eis}^k(\mathbf{x}) \cup p_2^* \text{Eis}^\ell(\mathbf{y}) \in K_{k+\ell+2}(E(N)^{k+\ell}) \otimes \mathbf{Q},$$

where $p_1 : E^{k+\ell} \rightarrow E^k$ and $p_2 : E^{k+\ell} \rightarrow E^\ell$ are the projections.

Conjecture

Let $k, \ell \geq 0$ and $\mathbf{x}, \mathbf{y}, \mathbf{z} \in (\mathbf{Z}/N\mathbf{Z})^2$ with $\mathbf{x} + \mathbf{y} + \mathbf{z} = 0$. Then

$$X^k Y^\ell(\mathbf{x}, \mathbf{y}) + X^\ell (-X - Y)^k(\mathbf{y}, \mathbf{z}) + Y^k (-X - Y)^\ell(\mathbf{z}, \mathbf{x}) = 0.$$

Conjecture

Let $k, \ell \geq 0$ and $\mathbf{x}, \mathbf{y}, \mathbf{z} \in (\mathbf{Z}/N\mathbf{Z})^2$ with $\mathbf{x} + \mathbf{y} + \mathbf{z} = 0$. Then

$$X^k Y^\ell(\mathbf{x}, \mathbf{y}) + X^\ell (-X - Y)^k(\mathbf{y}, \mathbf{z}) + Y^k (-X - Y)^\ell(\mathbf{z}, \mathbf{x}) = 0.$$

- ▶ One should be able to prove this in Deligne cohomology.
- ▶ Induction on the weight, using differentiation with respect to the parameters of the Eisenstein symbols.

Conjecture

Let $k, \ell \geq 0$ and $\mathbf{x}, \mathbf{y}, \mathbf{z} \in (\mathbf{Z}/N\mathbf{Z})^2$ with $\mathbf{x} + \mathbf{y} + \mathbf{z} = 0$. Then

$$X^k Y^\ell(\mathbf{x}, \mathbf{y}) + X^\ell (-X - Y)^k(\mathbf{y}, \mathbf{z}) + Y^k (-X - Y)^\ell(\mathbf{z}, \mathbf{x}) = 0.$$

- ▶ One should be able to prove this in Deligne cohomology.
- ▶ Induction on the weight, using differentiation with respect to the parameters of the Eisenstein symbols.
- ▶ Open question: what is the triangulation?

Conjecture

Let $k, \ell \geq 0$ and $\mathbf{x}, \mathbf{y}, \mathbf{z} \in (\mathbf{Z}/N\mathbf{Z})^2$ with $\mathbf{x} + \mathbf{y} + \mathbf{z} = 0$. Then

$$X^k Y^\ell(\mathbf{x}, \mathbf{y}) + X^\ell (-X - Y)^k(\mathbf{y}, \mathbf{z}) + Y^k (-X - Y)^\ell(\mathbf{z}, \mathbf{x}) = 0.$$

- ▶ One should be able to prove this in Deligne cohomology.
- ▶ Induction on the weight, using differentiation with respect to the parameters of the Eisenstein symbols.
- ▶ Open question: what is the triangulation?
- ▶ In this range, Deligne cohomology is just de Rham cohomology, so this amounts to say that a particular differential form is exact. Can we make explicit a primitive?

Beyond the reach of current technology

Conjecture (Rodriguez Villegas, 2003)

$$m(1 + x_1 + x_2 + x_3 + x_4) = -L'(f, -1)$$

$$m(1 + x_1 + x_2 + x_3 + x_4 + x_5) = -8L'(g, -1)$$

for modular forms $f \in S_3(\Gamma_1(15))$ and $g \in S_4(\Gamma_0(6))$.

Beyond the reach of current technology

Conjecture (Rodriguez Villegas, 2003)

$$m(1 + x_1 + x_2 + x_3 + x_4) = -L'(f, -1)$$

$$m(1 + x_1 + x_2 + x_3 + x_4 + x_5) = -8L'(g, -1)$$

for modular forms $f \in S_3(\Gamma_1(15))$ and $g \in S_4(\Gamma_0(6))$.

Conjecture (B.–Pengo, 2023)

$$m(xyt + xzt + yzt + xy + xz - yz - yt + zt - y + z - t + 1) = \frac{1}{6}L'(E, -2)$$

where $E = 32a2$ is an elliptic curve of conductor 32.

How we found the polynomial

Take $P(x, y, z, t)$ of the form

$$P = a(x, y) + b(x, y)z + c(x, y)t + d(x, y)zt.$$

Eliminating t in $P(x, y, z, t) = P(\frac{1}{x}, \frac{1}{y}, \frac{1}{z}, \frac{1}{t}) = 0$ gives

$$W_P : A(x, y)z^2 + B(x, y)z + C(x, y) = 0.$$

Want: $\Delta = B^2 - 4AC$ is a square $\delta(x, y)^2$ in $\mathbf{Q}(x, y)$.

Then $W_P = W_1 \cup W_2$ with

$$W_1 \cap W_2 : \delta(x, y) = 0.$$

We look for a, b, c, d such that $W_1 \cap W_2$ is an elliptic curve.

Numerical computation of $m(P)$

Rodriguez Villegas: $2m(P) = \log k - \int_0^{1/k} \phi_P(u) du$ where k is the constant coefficient of $P(x, y, z, t)P(\frac{1}{x}, \frac{1}{y}, \frac{1}{z}, \frac{1}{t})$ and

$$\phi_P(u) = \frac{1}{(2\pi i)^n} \int_{T^n} \frac{Q}{1-uQ} \cdot \frac{dx}{x} \frac{dy}{y} \frac{dz}{z} \frac{dt}{t}$$

with $Q = P(x, y, z, t)P(\frac{1}{x}, \frac{1}{y}, \frac{1}{z}, \frac{1}{t}) - k$.

Pengo–Ringeling: Using creative telescoping, one can find a polynomial ODE satisfied by ϕ_P . This takes a long time, but then $m(P)$ can be computed quickly with high precision.