

G. W. Gibbons

Carroll Symmetry, Gravitational Memory and Soft Gravitons
Black Holes, Quantum Information, Entanglement and all that

IHES

Bures sur Ivette Thursday 1 June 2017

May 31, 2017

Plan of Talk

The talk is based on work over the years with Christian Duval, Peter Horvathy and Pengming Zhang , much of it carried out at LMTP in Tours supported by a LE STUDIUM chair under the aegis of a collaborative project entitled: Classical and Quantum Space-Time and its Symmetries The talk falls into two parts.

- An introduction and overview of the Carroll Group
- An application to plane gravitational waves, gravitational memory and its relation to notion of a soft graviton

Contents of Part 1

- Galilei and Carroll groups as Kinematic groups obtained by contraction
- Isomorphism of Galilei and Carroll groups in $1 + 1$ spacetime dimensions
- Carrollian and Newton-Cartan Structures
- Bargmann Structures
- Carrollian and Newton-Cartan Structures obtained from a Bargmann structure in one higher dimension: Duality as projection versus pull back.
- Carrollian Automorphisms and Conformal Automorphisms
- Isomorphism of Conformal Carroll and Conformal Galilei algebras in $1 + 1$ spacetime dimensions
- BMS, and Newman-Unti groups as special cases
- Some Applications

The basic idea of our work is to look at **non-Einsteinian Relativity Principles** from an, albeit anachronistic, **Spacetime view point**

In our context a **Principle of Relativity** involves a notion of the **invariance of physical laws under passing to a moving frame** which we interpret as a **symmetry of some sort of spacetime structure**.

We follow the path pioneered by Bacry and Levy-Leblond * who found all algebras containing rotations, spatial and temporal translations and **boosts**. All may be regarded as Wigner-Inönü contractions † of the two De-Sitter groups.

Without boosts we would simply be classifying **Aristotelian spacetimes** which leads to Helmholtz's classification of congruence geometries ‡

.

*H. Bacry and J. Levy-Leblond, Possible kinematics J. Math. Phys. **9** (1968) 1605.

†E. Inönü , E.P. Wigner (1953). "On the Contraction of Groups and Their Representations". Proc. Nat. Acad. Sci. 39 (6): 51024.

‡Über die Thatsachen, welche der Geometrie zu Grunde liegen, in Wissenschaftliche Abhandlungen, Volume II, Leipzig: Johann Ambrosius Barth, 618639. Originally published in the Nachrichten von der Knigl. Gesellschaft der Wissenschaften zu Gttingen, No. 9 (3 June 1868).

The contractions are:

- Newton-Hooke $\Lambda \rightarrow 0, \quad c \rightarrow \infty, \quad c^2 \frac{\Lambda}{3}$ finite
- Poincaré $\Lambda \rightarrow 0, \quad c$ finite.
- Galilei $\Lambda \rightarrow 0, \quad c \rightarrow \infty.$
- Carroll $\Lambda \rightarrow 0, \quad c \rightarrow 0$

There is a certain **duality between the Galilei and Carroll groups**. In one the future light cone $t > \frac{1}{c}|\mathbf{x}|$ expands to become a future half space $t > 0$. In the other it contracts to become a future half line $t > 0, \mathbf{x} = 0$. One allows instantaneous propagation, the other is **ultra-local** and forbids any propagation.

All kinematic groups have a flat invariant **model space time** which allows a curved generalisation.

For Galilei this is Newton-Cartan spacetime with its degenerate co-metric g^{ij} whose kernel are co-normals of the absolute time slices

Carrollian spacetime. has a degenerate metric g_{ij} whose kernel is tangent to the absolute future *.

*To quote Mrs Thatcher: TINA, i.e. There is no alternative

Well, in our country," said Alice, still panting a little, "you'd generally get to somewhere else if you run very fast for a long time, as we've been doing."

A slow sort of country!" said the Queen. "Now, here, you see, it takes all the running you can do, to keep in the same place. If you want to get somewhere else, you must run at least twice as fast as that!"

Galilei, boosts act as

$$(t, \mathbf{x}) \rightarrow (t, \mathbf{x} - \mathbf{v}t)$$


Carroll, boosts act as

$$(s, \mathbf{x}) \rightarrow (s - \mathbf{b} \cdot \mathbf{x}, \mathbf{x})$$

where t is Galilean time and s is Carrollian time.

In 1+1 spacetime dimensions, Galileo and Carroll coincide as groups since we may interchange Galilean space and with Carrollian time and

vice versa



Taking the limit $c \uparrow \infty$ in the contra-variant Minkowski co- metric

$$-\frac{1}{c^2} \frac{\partial}{\partial t} \otimes \frac{\partial}{\partial t} + \delta^{ij} \frac{\partial}{\partial x^i} \frac{\partial}{\partial x^j}$$

motivates the definition of a **Newton-Cartan Spacetime** as a quadruple $\{N, \gamma, \theta, \nabla\}$ where N is a smooth $d+1$ manifold , γ a symmetric semi-positive definite contravariant 2-tensor of rank d with kernel the one-form θ and ∇ a symmetric affine connection w.r.t. which γ and θ are parallel.

Taking the limit $c \downarrow 0$ in the co-variant Minkowski metric

$$-c^2 dt^2 + \delta_{ij} dx^i dx^j$$

motivates the definition of a **Carrollian Spacetime** as a quadruple $\{C, g, \xi, \nabla\}$ where N is a smooth $d + 1$ manifold, g a symmetric semi-positive definite co-variant 2-tensor of rank d with kernel the vector field ξ and ∇ a symmetric affine connection w.r.t. which ξ and ∇ are parallel.

The standard flat case is $C = \mathbb{R} \times \mathbb{R}^d$, $g_{ij} = \delta_{ij}$, $\xi = \frac{\partial}{\partial s}$, $\Gamma_{\mu}{}^{\nu}{}_{\lambda} = 0$ where s is Carrollian time. The isometry group of the Carrollian metric contains

$$x^i \rightarrow x^i, \quad s \rightarrow s + f(x^i)$$

and so is infinite dimensional but if we require that the Carrollian automorphisms preserve the connection ∇ we obtain the standard finite dimensional Carroll group.

All the kinematic groups have a description in terms of Lorentzian geometry in 4+1 spacetime dimensions.

- Minkowski spacetime arises from a Kaluza-Klein reduction on a spacelike translation as shown by Kaluza and Klein.
- Newton-Cartan spacetime arises from a reduction on a null translation as shown by Duval et al. *
- Carrollian spacetime arises as the pull-back to a null hyperplane. Indeed given any null surface (like future null infinity \mathcal{I}^+) Carrollian structures come into play.

*C. Duval, G. Burdet, H. P. Kunzle and M. Perrin, Bargmann Structures and Newton-cartan Theory Phys. Rev. D **31** (1985) 1841. doi:10.1103/PhysRevD.31.1841

There is a fascinating link here with Cohen and Glashow's **Very Special Relativity***. They suggest that CP violation is so weak because the Lorentz group $S(3, 1)$ is weakly broken to its largest dimensional subgroup, $SIM(2)$, together with translations to make $ISM(2)$, which is 8 dimensional. It is the subgroup of $ISO(3, 1)$ which preserves a null direction and may be considered the Bargman group of $2 + 1$ dimensional Newton-Cartan spacetime.

*A. G. Cohen and S. L. Glashow, Very special relativity Phys. Rev. Lett. **97** (2006) 021601

$$ds^2 = -2dudv + dx^i dx^i \quad i = 1, \dots, n-2$$

$$p_i = \partial_i, \quad L_{ij} = x_i \partial_j - x_j \partial_i$$

$$U = \partial_u, \quad V = \partial_v, \quad N = u \partial_i - v \partial_v$$

$$U_i = u \partial_i + x_i \partial_v, \quad V_i \partial_i + x_i \partial_v$$

$$\text{Bargmann}(n-2, 1) : \quad , \quad \text{span}\{p_i, L_{ij}, U, V, U_i\}$$

$$\text{Gailili}(n-2, 1) : \quad \text{Bargmann}(n, 1)/V$$

$$\text{Carroll}(n-2, 1) : \quad \text{span}\{p_i, L_{ij}, V, U_i\}$$

$$\text{Sim}(n-2) : \quad \text{span}\{U_i, L_{ij}, N\}$$

$$\text{ISim}(n-2) : \quad \text{span}\{p_i, U, V, U_i, L_{ij}, N\}$$

We define a **Bargmann Manifold** as a triple $\{B, G, \xi\}$ where B is a $(d+2)$ manifold, G a Lorentzian metric (i.e non-degenerate and signature $(d+1, 1)$) and a null vector field ξ which is parallel w.r.t. the Levi-Civita connection of G . The standard flat Bargmann structure is given by $B = \mathbb{R}^{d+2}$, $\xi = \frac{\partial}{\partial s}$ with

$$ds^2 = \delta_{ij} dx^i dx^j + 2dt ds$$

Note that *both* s and t are null coordinates.

The standard flat Newton-Cartan structure is obtained by **pushing forward** the flat Bargmann structure to the quotient or **lightlike shadow** or **null reduction** $N = B/(\mathbb{R}\xi)$. The Bargmann group consists of those isometries of B which preserve ξ . This is a central extension of the Galilei group, the centre being generated by ξ .

One may also obtain the central extension of the conformal Schrödinger group, the symmetry of the free Schrödinger equation as the those conformal transformations of $d + 2$ -dimensional Minkowski spacetime which commute with the action of $\mathbb{R}\xi$.

A massless scalar field in $\mathbb{E}^{d+1,1}$ is invariant under conformal transformations

$$2\frac{\partial^2\phi}{\partial t\partial s}\phi(s, t, x^i) + \nabla^2\phi = 0.$$

set

$$\xi\phi = -im\phi, \quad \phi = e^{-ims}\psi(t, x^i)$$

then

$$i\frac{\partial\psi}{\partial t} = -\frac{1}{2m}\nabla^2\psi.$$

The standard flat Carroll structure is obtained by **pulling back** the flat Bargmann structure to a **null hypersurface** $t = \text{constant}$. The Carroll group consists of these isometries of B which commute with the pull back.

By a Lie-algebra co-homology argument it has been shown that that the Carroll group admits no central extension.

A non-standard Carroll structure may be obtained by taking the product $C = \mathbb{R} \times \Sigma_d$ where Σ_d with Riemannian metric \hat{g} and $g = \hat{g} \oplus 0 \times du^2$ and $\xi = \frac{\partial}{\partial u}$, where u is a coordinate on \mathbb{R} . For ∇ we *could* take the Levi-civita connection of $\{\Sigma, \hat{g}\}$.

For a general Carroll structure $\{C, g, \xi \nabla\}$ we define the **Conformal Carroll group of level N** as consisting of diffeomorphisms a such that

$$a^* \hat{g} = \Omega^2 \hat{g}, \quad a_* = \Omega^{-\frac{2}{N}} \xi$$

For the flat Carroll structure this has Killing vector fields

$$X = (\omega_{ij} x_j + \gamma_i (\chi - 2\kappa_i x_i) + \kappa_i x_j x_j) \frac{\partial}{\partial x_i} + \left(\frac{2}{N} (\chi - 2\kappa_j x_j) u + T(x_k) \right) \frac{\partial}{\partial u}$$

This is *infinite dimensional* because of the **super-translations** $T(x_i)$ which have conformal weight $= -\frac{2}{N}$, i.e. are densities of weight $\nu = -\frac{2}{Nd}$. The quantity $z = \frac{2}{N}$ is known as a **dynamical exponent**.

If $N = 2$, $z = 1$ and we have symmetry between the scaling of space and time.

If $d = 1$, using the isomorphism between the Carroll and Galilei algebras described above we obtain the **Conformal Galilei algebra**. **CGA** introduced by many people in a variety of contexts.

The **isometry group of the flat Carroll structure** is obtained by setting $\Omega = 1$. Its Lie algebra is also infinite dimensional, because of the supertanslations. Requiring that the connection is preserved reduces the Carroll Lie algebra to the standard finite dimensional case obtained by Levy-Lebond and Bacry.

For a general Carrollian structure , the Conformal Carroll group is generated by

$$X = Y + \left(\frac{\lambda}{N} + T(x) \right) \frac{\partial}{\partial u}$$

where Y is a conformal vector field of $\{\Sigma, \hat{g}\}$

$$\mathcal{L}_Y g = \lambda g$$

generating

$$x \rightarrow \phi(x), \quad u \rightarrow \Omega^{\frac{2}{N}}(u + \alpha(x))$$

Example

If $\{\Sigma_d, \hat{g}\} = \{S^1, d\theta^2\}$ we get $\text{Diff}(S^1)$ semi-direct product super translations of weight $\nu = -\frac{2}{N}$ generated by the vector field

$$X = Y(\theta)\frac{\partial}{\partial\theta} + \left(\frac{2}{N}Y'(\theta) + T(\theta)\right)\frac{\partial}{\partial u}.$$

whose algebra is an extension of the **Witt** or **Virasoro** algebra.

Example

If $\{\Sigma_d, g\} = \{S^2, d\theta^2 + \sin^2\theta d\phi^2\}$ and $N = 2$ we get

$$PSL(2, \mathbb{C}) \ltimes \mathcal{T}$$

where \mathcal{T} are half densities on S^2 which is the **Bondi-Metzner-Sachs Group**

Which was originally discovered as the asymptotic symmetry group of an asymptotically flat four-dimensional spacetime. The BMS Group has an obvious generalisation to S^d for all $d > 2$. However this generalisation does not appear to coincide with the asymptotic symmetry group of an asymptotically flat spacetime of dimension greater than four.

We may weaken our requirement on the conformal Carroll group so that $\alpha(x)$ only preserves the conformal class of the Carroll metric g . Since the vector field ξ spans the kernel of g , α is only required to preserve the *direction* of ξ . Thus its generating vector field X need only satisfy

$$\mathcal{L}_X \xi = \mu(x, u) \xi, \quad \Rightarrow X = Y + F(u, x) \frac{\partial}{\partial u} \quad (1)$$

where $F(u, x)$ is an arbitrary function of *both* u and x , and Y is a conformal Killing field of the Riemannian manifold $\{\Sigma_d, \hat{g}\}$

Example

If $\{\Sigma_d, \hat{g}\} = \{\Sigma_d, g\} = \{S^2, d\theta^2 + \sin^2 \theta d\phi^2\}$ we obtain the **Newman-Unti Group** which was introduced by these authors in the study of asymptotically flat four-dimensional spacetimes.

- Carrollian and BMS symmetries have a number of applications to various topics of current interest to string theorists and holographers which was part of the original motivation for the work reported here.
- Carrollian symmetries arise in **tachyon condensates** and the strong coupling limits of Born-Infeld theories on brane. * .
- Using our enhanced understanding of the Carroll group we were able to construct Carrollian-invariant theories of electromagnetism which potentially have applications to **slow light**.
- Using a geometric quantization method of Souriau we constructed theories of **Carrollian massive and massless particles**. One finds the former do not move, consistent with other view points.

*G. Gibbons, K. Hashimoto and P. Yi, "Tachyon condensates, Carrollian contraction of Lorentz group, and fundamental strings," JHEP **0209** (2002) 061

- One of the most intriguing was to **Schild or Null Strings**, that is strings whose two-dimensional world sheet carries a Carrollian metric, i.e. is a two-dimensional null surface. It turns out that Souriau's procedure for obtaining dynamical systems invariant under a group G applied to massless "particles " leads to Schild Strings.

Our most recent application has been to the **Carroll Symmetry** of **Plane gravitational waves** and to **Soft Gravitons & the Memory Effect for Plane Gravitational waves**.

These are vacuum metrics with a covariantly constant null Killing vector field and they admit a five-dimensional isometry group acting on null hypersurfaces and contains a three-dimensional abelian subgroup.

The isometry group is a 5-dimensional subgroup of the 6 dimensional Carroll group in 2+1 dimensions with the rotations broken

For a scalar wave it would be the full Carroll group

These solutions have many remarkable properties.

- They admit a covariantly constant spinor field. They are thus BPS.
- All invariants constructed from the Riemann tensor and its covariant derivatives
- They suffer no quantum corrections
- They are exact solutions of string theory

These metrics admit two useful coordinates systems.

Brinkmann coordinates

$$ds^2 = \delta_{ij} dX^i dX^j + 2dU dV + K_{ij}(U) X^i X^j dU^2, \quad \text{Tr}K = 0.$$

Baldwin-Jeffery-Rosen Coordinates

$$ds^2 = a_{ij}(u) dx^i dx^j + 2du dv$$

$$\mathbf{X} = P(u)\mathbf{x}, \quad U = u, \quad V = v - \frac{1}{4}\mathbf{x} \cdot a(u)\mathbf{x}$$

$$a = P^t P, \quad \ddot{P} = KP, \quad P^t \dot{P} - \dot{P}^t P = 0.$$

$$K = \frac{1}{2}P\left(\dot{b} + \frac{1}{2}b^2\right)P^{-1}, \quad b = a^{-1}\dot{a}.$$

In Brinkmann coordinates the field equations are trivially satisfied


$$K_{11} = -K_{22} = \frac{1}{2}\mathcal{A}_+(U), \quad K_{12} = K_{21} = \frac{1}{2}\mathcal{A}_\times(U)$$

where the two polarization amplitudes may be given as **arbitrary** functions of U .

In Baldwin-Jeffery-Rosen coordinates the field equations are highly non-linear and even the flat solutions are non-trivial. However the high degree of manifest symmetry in BJR coordinates **allows exact solution of all geodesics**



Consider a sandwich wave*



passing over a cloud of particles all at relative rest before the wave arrives.

*for which $\mathcal{A}_+(U)$ and $\mathcal{A}_\times(U)$ vanish outside a finite interval $U_i \leq U \leq U_f$.

By Noether's theorem the Baldwin-Jeffery-Rosen coordinates x are constant both before and after the wave has passed

But even if $a_{ij} = \delta_{ij}$ before the wave has arrived a_{ij} will, in general have non-trivial time dependence after the wave has passed. Thus the separations of the particles will in general have non-trivial time dependence after the wave has passed. This may in principle be measured and information about $\mathcal{A}_+(U)$ and $\mathcal{A}_\times(U)$ deduced.

THIS IS THE GRAVITATIONAL MEMORY EFFECT

A **particular case**, whose existence was first explicitly pointed out by Zeldovich and Polnarev (Ya. B. Zel'dovich and A. G. Polnarev, Radiation of gravitational waves by a cluster of superdense stars, Sov. Astron. **18** 17 (1974)) is when $a_{ij}(U_i) = \delta_{ij}$ and $a_{ij}(U_f) \neq \delta_{ij}$ but $\dot{a}_{ij}(U_f) = 0$ which leads to a permanent constant separation different from the initial separation whose existence was implicit in earlier work (G. W. Gibbons and S. W. Hawking, Theory of the detection of short bursts of gravitational radiation, Phys. Rev. D **4** (1971) 219).



This is sometimes crassly referred to as “a permanent change of spacetime after the wave has passed.”. It is no such thing.

It is known from the work of Tamm and Plebanski that for a metric of the form

$$ds^2 = -dt^2 + g(x, t)_{ab} dx^a dx^b$$

solutions of Maxwell's equations behave as if in an **impedance matched** medium in flat spacetime with

$$\epsilon_{ab} = \mu_{ab} = \sqrt{\det g} (g^{-1})_{ab}$$



In BJR coordinates

$$\epsilon_{ij} = \sqrt{\det a} (a^{-1})_{ij}, \quad \epsilon_{33} = \sqrt{\det a}$$

After the wave has passed although $a_{ij} \neq \delta_{ij}$ there is a coordinate transformation which we calculate explicitly, which brings the metric after the wave has passed to canonical flat form. **This coordinate transformation does not tend to the identity at spatial infinity.**

THUS FLAT PLANE WAVE VACCUUM METRICS IN BALDWIN-JEFFERY-ROSEN COORDINATES CAN BE THOUGHT OF AS SOFT GRAVITONS LEFT AFTER THE PASSING OF A WAVE PULSE WITH NON-VANISHING CURVATURE

This talk was based on the following papers

P. M. Zhang, C. Duval, G. W. Gibbons and P. A. Horvathy, Soft Gravitons and the Memory Effect for Plane Gravitational Waves [arXiv:1706.01378 \[gr-qc\]](#).

P. M. Zhang, C. Duval, G. W. Gibbons and P. A. Horvathy, The Memory Effect for Plane Gravitational Waves, [arXiv:1704.05997 \[gr-qc\]](#).

C. Duval, G. W. Gibbons, P. A. Horvathy and P.-M. Zhang, Carroll symmetry of plane gravitational waves," [arXiv:1702.08284 \[gr-qc\]](#).

C. Duval, G. W. Gibbons and P. A. Horvathy Conformal Carroll groups, *J. Phys. A* **47** (2014) 335204 [[arXiv:1403.4213 \[hep-th\]](#)]

C. Duval, G. W. Gibbons and P. A. Horvathy, 'Conformal Carroll groups and BMS symmetry,' *Class. Quant. Grav.* **31** (2014) 092001 [[arXiv:1402.5894 \[gr-qc\]](#)]

C. Duval, G. W. Gibbons, P. A. Horvathy and P. M. Zhang, 'Carroll versus Newton and Galilei: two dual non-Einsteinian concepts of time, *Class. Quant. Grav.* **31** (2014) 085016 [[arXiv:1402.0657 \[gr-qc\]](#)]

C. Duval, G. W. Gibbons and P. Horvathy, Celestial mechanics, conformal structures and gravitational waves, *Phys. Rev. D* **43** (1991) 3907 [[hep-th/0512188](#)].