

Integrable Lagrangians and Picard modular forms

E.V. Ferapontov

Department of Mathematical Sciences, Loughborough University, UK

E.V.Ferapontov@lboro.ac.uk

Based on ongoing project with F. Cléry, A. Odesskii, D. Zagier

Classical and Quantum Integrability

2-6 September 2019, Dijon

Plan:

- Integrable Lagrangians $\int f(v_{x_1}, v_{x_2}, v_{x_3}) dx_1 dx_2 dx_3$
- Integrability conditions
- Lagrangian densities $f = v_{x_1} v_{x_2} g(v_{x_3})$
- Lagrangian densities $f = v_{x_1} g(v_{x_2}, v_{x_3})$
- Generic Lagrangian densities $f(v_{x_1}, v_{x_2}, v_{x_3})$

References:

- [1] E.V. Ferapontov, K.R. Khusnutdinova and S.P. Tsarev, On a class of three-dimensional integrable Lagrangians, Comm. Math. Phys. **261**, N1 (2006) 225-243.
- [2] E.V. Ferapontov and A.V. Odesskii, Integrable Lagrangians and modular forms, J. Geom. Phys. **60**, no. 6-8 (2010) 896-906.
- [3] D. Zagier, On a U(3,1)-automorphic form of Ferapontov-Odesskii, talk in Utrecht on 17 April 2009.
- [4] A.V. Odesskii, V.V. Sokolov, Integrable pseudopotentials related to generalized hypergeometric functions, Selecta Math. **16** (2010) 145-172.

Integrable Lagrangians $\int f(v_{x_1}, v_{x_2}, v_{x_3}) dx_1 dx_2 dx_3$

Euler-Lagrange equation:

$$(f_{v_{x_1}})_{x_1} + (f_{v_{x_2}})_{x_2} + (f_{v_{x_3}})_{x_3} = 0.$$

Examples:

Dispersionless Kadomtsev-Petviashvili (dKP) equation

$$v_{x_1 x_3} - v_{x_1} v_{x_1 x_1} - v_{x_2 x_2} = 0, \quad f = v_{x_1} v_{x_2} - \frac{1}{3} v_{x_1}^3 - v_{x_2}^2.$$

Boyer-Finley (BF) equation

$$v_{x_1 x_1} + v_{x_2 x_2} - e^{v_{x_3}} v_{x_3 x_3} = 0, \quad f = v_{x_1}^2 + v_{x_2}^2 - 2e^{v_{x_3}}.$$

Main problem

- The parameter space of integrable Lagrangian densities f is 20-dimensional.
- Integrability conditions for f are invariant under a 20-dimensional symmetry group which acts on the parameter space with an open orbit.

Question: What is the **master-Lagrangian** corresponding to the open orbit?

We will see that it is related to the theory of Picard modular forms.

Three equivalent approaches to integrability

- The method of hydrodynamic reductions based on the requirement that the equation possesses infinitely many multi-phase solutions of special type.
- The method of dispersionless Lax pairs based on the representation of the equation as the compatibility condition of two Hamilton-Jacobi type equations.
- Integrability 'on solutions' based on the condition that the characteristic variety of the equation defines a conformal structure which is Einstein-Weyl on every solution.

All three approaches lead to the same set of integrability conditions for the Lagrangian density f .

Hydrodynamic reductions: example of dKP

First-order form of dKP equation $v_{x_1 x_3} - v_{x_1} v_{x_1 x_1} - v_{x_2 x_2} = 0$ (set $u = v_{x_1}$):

$$u_{x_3} - u u_{x_1} - w_{x_2} = 0, \quad u_{x_2} - w_{x_1} = 0.$$

Look for N -phase solutions: $u = u(R^1, \dots, R^n)$, $w = w(R^1, \dots, R^n)$ where

$$R_{x_3}^i = \lambda^i(R) R_{x_1}^i, \quad R_{x_2}^i = \mu^i(R) R_{x_1}^i.$$

The substitution of u, w into the above first-order system implies

$$\partial_i w = \mu^i \partial_i u, \quad \lambda^i = u + (\mu^i)^2,$$

as well as the following equations for $u(R)$ and $\mu^i(R)$ (Gibbons-Tsarev system):

$$\partial_j \mu^i = \frac{\partial_j u}{\mu^j - \mu^i}, \quad \partial_i \partial_j u = 2 \frac{\partial_i u \partial_j u}{(\mu^j - \mu^i)^2}.$$

In involution! General solution depends on n arbitrary functions of one variable.

J. Gibbons, S.P. Tsarev, Reductions of the Benney equations, Phys. Lett. A **211** (1996) 19-24.

Dispersionless Lax pairs: example of dKP

The dKP equation $v_{x_1 x_3} - v_{x_1} v_{x_1 x_1} - v_{x_2 x_2} = 0$ possesses dispersionless Lax representation

$$S_{x_2} = \frac{1}{2} S_{x_1}^2 + v_{x_1}, \quad S_{x_3} = \frac{1}{3} S_{x_1}^3 + v_{x_1} S_{x_1} + v_{x_2}.$$

In parametric form:

$$S_{x_1} = p, \quad S_{x_2} = \frac{1}{2} p^2 + v_{x_1}, \quad S_{x_3} = \frac{1}{3} p^3 + v_{x_1} p + v_{x_2}.$$

E.V. Zakharov, Dispersionless limit of integrable systems in $2 + 1$ dimensions, in: Singular Limits of Dispersive Waves, Ed. N.M. Ercolani et al., Plenum Press, NY (1994) 165-174.

Observation Integrability by the method of hydrodynamic reductions is equivalent to the existence of a dispersionless Lax representation.

Integrability via Einstein-Weyl geometry: example of dKP

Einstein-Weyl geometry is a triple (\mathbb{D}, g, ω) where \mathbb{D} is a symmetric connection, g is a conformal structure and ω is a covector such that

$$\mathbb{D}_k g_{ij} = \omega_k g_{ij}, \quad R_{(ij)} = \Lambda g_{ij}.$$

Here $R_{(ij)}$ is the symmetrised Ricci tensor of \mathbb{D} and Λ is some function (the first set of equations defines \mathbb{D} uniquely, so it is sufficient to specify g and ω only).

Every solution of the dKP equation $v_{x_1 x_3} - v_{x_1} v_{x_1 x_1} - v_{x_2 x_2} = 0$ carries Einstein-Weyl geometry (Dunajski, Mason, Tod):

$$g = 4dx_1 dx_3 - dx_2^2 + 4v_{x_1} dx_3^2, \quad \omega = -4v_{x_1 x_1} dx_3.$$

Observation Integrability is equivalent to the Einstein-Weyl property of the characteristic conformal structure of the equation.

E.V. Ferapontov, B.S. Kruglikov, Dispersionless integrable systems in 3D and Einstein-Weyl geometry, J. Diff. Geom. **97** (2014) 215-254.

Integrability conditions

For a non-degenerate Lagrangian, the Euler-Lagrange equation is integrable (by either of the techniques mentioned above) if and only if the Lagrangian density f satisfies the relation

$$d^4 f = d^3 f \frac{dH}{H} + \frac{3}{H} \det(dM).$$

Here $d^3 f$ and $d^4 f$ are the symmetric differentials of f while the Hessian H and the 4×4 augmented Hessian matrix M are defined as

$$H = \det \begin{pmatrix} f_{xx} & f_{xy} & f_{xz} \\ f_{xy} & f_{yy} & f_{yz} \\ f_{xz} & f_{yz} & f_{zz} \end{pmatrix}, \quad M = \begin{pmatrix} 0 & f_x & f_y & f_z \\ f_x & f_{xx} & f_{xy} & f_{xz} \\ f_y & f_{xy} & f_{yy} & f_{yz} \\ f_z & f_{xz} & f_{yz} & f_{zz} \end{pmatrix}.$$

Here $(x, y, z) = (v_{x_1}, v_{x_2}, v_{x_3})$. The non-degeneracy condition is equivalent to $H \neq 0$. The system for f is in involution, and its solution space is 20-dimensional.

Weierstrass sigma function σ and integers B_k

Let σ be the Weierstrass sigma function of the elliptic curve $y^2 = 4x^3 - \frac{1}{2}$ (case $g_2 = 0, g_3 = \frac{1}{2}$). This function satisfies the fourth-order ODE

$$\sigma\sigma'''' - 4\sigma'\sigma'' + 3(\sigma'')^2 = 0$$

and possesses a power series expansion

$$\sigma(z) = \sum_{k \geq 0} B_k \frac{z^{6k+1}}{(6k+1)!}$$

where B_k are certain integers.

Lagrangian density $f = v_{x_1} v_{x_2} v_{x_3}$

Euler-Lagrange equation:

$$v_{x_3} v_{x_1 x_2} + v_{x_2} v_{x_1 x_3} + v_{x_1} v_{x_2 x_3} = 0.$$

Parametric Lax representation:

$$\frac{S_{x_1}}{v_{x_1}} = \zeta(p) + \frac{\wp'(p) + \lambda}{2\wp(p)}, \quad \frac{S_{x_2}}{v_{x_2}} = \zeta(p) + \frac{\wp'(p) - \lambda}{2\wp(p)}, \quad \frac{S_{x_3}}{v_{x_3}} = \zeta(p),$$

where $(\wp')^2 = 4\wp^3 + \lambda^2$ and $\zeta' = -\wp$ (Weierstrass \wp and ζ functions).

Lagrangian densities $f = v_{x_1} v_{x_2} g(v_{x_3})$

Euler-Lagrange equation:

$$(v_{x_2} g(v_{x_3}))_{x_1} + (v_{x_1} g(v_{x_3}))_{x_2} + (v_{x_1} v_{x_2} g'(v_{x_3}))_{x_3} = 0.$$

Integrability condition for $g(z)$:

$$g''''(g^2 g'' - 2g(g')^2) - 9(g')^2 (g'')^2 + 2gg'g''g''' + 8(g')^3 g''' - g^2 (g''')^2 = 0.$$

$GL(2, \mathbb{R})$ -invariance:

$$\tilde{z} = \frac{\alpha z + \beta}{\gamma z + \delta}, \quad \tilde{g} = (\gamma z + \delta)g.$$

This invariance allows one to linearise the integrability condition for $g(z)$.

Auxiliary hypergeometric equation

Consider the auxiliary hypergeometric equation

$$u(1-u)h_{uu} + (1-2u)h_u - \frac{2}{9}h = 0,$$

parameters $(1/3, 2/3, 1)$. The geometry behind this equation is a 1-parameter family of genus 2 trigonal curves

$$r^3 = q(q-1)(q-u)^2$$

supplied with the holomorphic differential $\omega = dq/r$. The corresponding periods, $h = \int_a^b \omega$ where $a, b \in \{0, 1, \infty, u\}$, form a 2-dimensional vector space and satisfy the above (Picard-Fuchs) hypergeometric equation.

Generic solution $g(z)$

The generic solution $g(z)$ can be represented in any of the three equivalent forms:

1. Parametric form:

$$z = \frac{h_1(u)}{h_2(u)}, \quad g = h_2(u)$$

where h_i are two linearly independent solutions of the hypergeometric equation. $GL(2, \mathbb{R})$ -invariance corresponds to the freedom in the choice of basis h_i .

2. Theta representation:

$$g(z) = \sum_{(k,l) \in \mathbb{Z}^2} e^{2\pi i(k^2 + kl + l^2)z} = 1 + 6q + 6q^3 + 6q^4 + 12q^4 + \dots, \quad q = e^{2\pi iz}.$$

3. Power series:

$$g(z) = \sum_{k \geq 0} B_k^2 \frac{z^{6k+1}}{(6k+1)!}$$

where B_k are integers (same as in the expansion of σ function).

Lagrangian densities $f = v_{x_1} g(v_{x_2}, v_{x_3})$

Euler-Lagrange equation

$$(g)_{x_1} + (v_{x_1} g_{v_{x_2}})_{x_2} + (v_{x_1} g_{v_{x_3}})_{x_3} = 0.$$

Integrability conditions lead to an involutive system of five PDEs for $g(y, z)$ which are invariant under the 10-dimensional symmetry group:

$$\tilde{y} = \frac{l_1(y, z)}{l(y, z)}, \quad \tilde{z} = \frac{l_2(y, z)}{l(y, z)}, \quad \tilde{g} = \alpha g + \beta,$$

where l, l_1, l_2 are arbitrary (inhomogeneous) linear forms. This invariance allows one to linearise the integrability conditions for $g(y, z)$.

Auxiliary hypergeometric system

Consider the auxiliary (Appell) hypergeometric system

$$h_{u_1 u_2} = \frac{1}{3} \frac{h_{u_1} - h_{u_2}}{u_1 - u_2},$$

$$h_{u_1 u_1} = -\frac{h}{9u_1(u_1 - 1)} + \frac{h_{u_2}}{3(u_1 - u_2)} \frac{u_2(u_2 - 1)}{u_1(u_1 - 1)} - \frac{h_{u_1}}{3} \left(\frac{1}{u_1 - u_2} + \frac{2}{u_1} + \frac{2}{u_1 - 1} \right),$$

$$h_{u_2 u_2} = -\frac{h}{9u_2(u_2 - 1)} + \frac{h_{u_1}}{3(u_2 - u_1)} \frac{u_1(u_1 - 1)}{u_2(u_2 - 1)} - \frac{h_{u_2}}{3} \left(\frac{1}{u_2 - u_1} + \frac{2}{u_2} + \frac{2}{u_2 - 1} \right).$$

The geometry behind this system is the family of genus 3 Picard trigonal curves

$$r^3 = q(q - 1)(q - u_1)(q - u_2)$$

supplied with the holomorphic differential $\omega = dq/r$. The corresponding periods,

$h = \int_a^b \omega$ where $a, b \in \{0, 1, \infty, u_1, u_2\}$, form a 3-dimensional vector space and satisfy the above (Picard-Fuchs) hypergeometric system.

Generic solution $g(y, z)$

The generic solution $g(y, z)$ can be represented in any of the 3 equivalent forms:

1. Parametric form:

$$y = \frac{h_1(u_1, u_2)}{h_3(u_1, u_2)}, \quad z = \frac{h_2(u_1, u_2)}{h_3(u_1, u_2)}, \quad g = F(s), \quad s = \frac{u_1(u_2 - 1)}{u_2(u_1 - 1)}$$

where h_i are three linearly independent solutions of the hypergeometric system and $F' = [s(s - 1)]^{-2/3}$.

2. Theta representation:

$$g(y, z) = y + \sum_{(k,l) \in \mathbb{Z}^2 \setminus 0} \frac{\sigma((k + \epsilon l)y)}{k + \epsilon l} e^{2\pi i(k^2 + kl + l^2)z}, \quad \epsilon = e^{\pi i/3}.$$

3. Power series:

$$g(y, z) = \sum_{j,k \geq 0} B_j B_k B_{j+k} \frac{y^{6j+1}}{(6j+1)!} \frac{z^{6k+1}}{(6k+1)!}.$$

Relation to Picard modular forms

The period map

$$y = \frac{h_1(u_1, u_2)}{h_3(u_1, u_2)}, \quad z = \frac{h_2(u_1, u_2)}{h_3(u_1, u_2)},$$

was inverted by Picard (1883):

$$u_1 = \frac{\varphi_1(y, z)}{\varphi_0(y, z)}, \quad u_2 = \frac{\varphi_2(y, z)}{\varphi_0(y, z)},$$

where φ_ν are single-valued modular forms on a 2-dimensional complex ball

$2\operatorname{Re}y + |z|^2 < 0$ with respect to the Picard modular group

$\Gamma[\sqrt{-3}] = \{g \in U(2, 1; \mathbb{Z}[\rho]) : g \equiv 1 \pmod{\sqrt{-3}}\}$, $\rho = e^{2\pi i/3}$. Picard

modular forms were extensively studied by Holzapfel, Feustel, Finis, Shiga, Cléry

and van der Geer.

Differential dg via Picard modular forms

There is a simple expression of the differential dg in terms of φ_ν :

$$dg = \frac{\varphi_1\varphi_2(\varphi_2 - \varphi_1)d\varphi_0 + \varphi_0\varphi_2(\varphi_0 - \varphi_2)d\varphi_1 + \varphi_0\varphi_1(\varphi_1 - \varphi_0)d\varphi_2}{\zeta^2}$$

where ζ is a modular form defined as

$$\zeta^3 = \varphi_0\varphi_1\varphi_2(\varphi_1 - \varphi_0)(\varphi_2 - \varphi_0)(\varphi_2 - \varphi_1).$$

Up to a constant factor, the differential dg coincides with the Eisenstein series $E_{1,1}$ which was known before from the theory of vector-valued Picard modular forms.

H. Shiga, On the representation of the Picard modular function by θ constants, I, II. Publ. Res. Inst. Math. Sci. **24**, no. 3 (1988) 311-360.

F. Cléry, G. van der Geer, Generators for modules of vector-valued Picard modular forms, Nagoya Math. J. **212** (2013) 19-57.

Generic Lagrangian densities $f(v_{x_1}, v_{x_2}, v_{x_3})$

Euler-Lagrange equation:

$$(f_{v_{x_1}})_{x_1} + (f_{v_{x_2}})_{x_2} + (f_{v_{x_3}})_{x_3} = 0.$$

Integrability conditions lead to a system of fifteen PDEs for $f(x, y, z)$ which are invariant under 20-dimensional symmetry group:

$$\tilde{x} = \frac{l_1(x, y, z)}{l(x, y, z)}, \quad \tilde{y} = \frac{l_2(x, y, z)}{l(x, y, z)}, \quad \tilde{z} = \frac{l_3(x, y, z)}{l(x, y, z)}, \quad \tilde{f} = \frac{f}{l(x, y, z)},$$

as well as obvious symmetries of the form

$$\tilde{f} = \epsilon f + \alpha x + \beta y + \gamma z + \delta,$$

where l, l_1, l_2, l_3 are arbitrary (inhomogeneous) linear forms. This invariance allows one to linearise the integrability conditions for $f(x, y, z)$.

Auxiliary hypergeometric system

Consider the auxiliary (Appell) hypergeometric system

$$h_{u_i u_j} = \frac{1}{3} \frac{h_{u_i} - h_{u_j}}{u_i - u_j},$$

$$h_{u_i u_i} = -\frac{2}{9} \frac{h}{u_i(u_i - 1)} - \frac{1}{3u_i(u_i - 1)} \sum_{j \neq i}^3 \frac{u_j(u_j - 1)}{u_j - u_i} h_{u_j} +$$

$$-\frac{1}{3} \left(\sum_{j \neq i}^3 \frac{1}{u_i - u_j} + \frac{2}{u_i} + \frac{2}{u_i - 1} \right) h_{u_i}.$$

The geometry behind this system is the family of genus 4 Picard trigonal curves

$$r^3 = q(q - 1)(q - u_1)(q - u_2)(q - u_3)$$

supplied with the holomorphic differential $\omega = dq/r$. The corresponding periods, $h = \int_a^b \omega$ where $a, b \in \{0, 1, \infty, u_1, u_2, u_3\}$, form a 4-dimensional vector space and satisfy the above (Picard-Fuchs) hypergeometric system.

Inhomogeneous hypergeometric extension

We will also need the inhomogeneous hypergeometric system

$$F_{u_i u_j} = \frac{1}{3} \frac{F_{u_i} - F_{u_j}}{u_i - u_j} + \epsilon_{ijk} \frac{u_k(u_k - 1)(u_i - u_j)}{U^{2/3}},$$

$$F_{u_i u_i} = -\frac{2}{9} \frac{F}{u_i(u_i - 1)} - \frac{1}{3u_i(u_i - 1)} \sum_{j \neq i}^3 \frac{u_j(u_j - 1)}{u_j - u_i} F_{u_j} + \\ -\frac{1}{3} \left(\sum_{j \neq i}^3 \frac{1}{u_i - u_j} + \frac{2}{u_i} + \frac{2}{u_i - 1} \right) F_{u_i},$$

where ϵ_{ijk} is the totally antisymmetric tensor and

$$U = u_1 u_2 u_3 (u_1 - 1)(u_2 - 1)(u_3 - 1)(u_1 - u_2)(u_2 - u_3)(u_3 - u_1).$$

The inhomogeneous system for F is in involution.

Generic solution $f(x, y, z)$

The generic solution $f(x, y, z)$ can be represented in any of the 3 equivalent forms:

1. Parametric form:

$$x = \frac{h_1}{h_4}, \quad y = \frac{h_2}{h_4}, \quad z = \frac{h_3}{h_4}, \quad f = \frac{F}{h_4}$$

where $h_i(u_1, u_2, u_3)$ are four independent solutions of the hypergeometric system and $F(u_1, u_2, u_3)$ is a solution of the inhomogeneous system.

2. Theta representation:

$$f(x, y, z) = xy + \sum_{(k,l) \in \mathbb{Z}^2 \setminus 0} \frac{\sigma((k + \epsilon l)x)\sigma((k + \epsilon l)y)}{(k + \epsilon l)^2} e^{2\pi i(k^2 + kl + l^2)z}.$$

3. Power series:

$$f(x, y, z) = \sum_{i,j,k \geq 0} B_i B_j B_k B_{i+j+k} \frac{x^{6i+1}}{(6i+1)!} \frac{y^{6j+1}}{(6j+1)!} \frac{z^{6k+1}}{(6k+1)!}.$$

Concluding remarks

- There exists integrable Lagrangians whose densities f are polynomial, or can be expressed in terms of elementary functions. It would be interesting to clarify how these (and similar) examples can be recovered as degenerations of the ‘master-Lagrangian’, and to describe singular orbits of lower dimension. This should be related to understanding degenerations/compactifications of the moduli space of Picard curves.
- Although our parametric, theta and power series representations for integrable densities possess straightforward generalisations to dimensions higher than 3, the relation to integrable Lagrangians will be lost: one can show that in higher dimensions every integrable first-order Lagrangian density $f(v_x)$ is necessarily of the form $f = \frac{Q(v_x)}{l(v_x)}$ where Q and l are arbitrary quadratic and linear functions of the first-order derivatives. Thus, the occurrence of modular forms is the essentially 3-dimensional phenomenon.