

Multimessenger opportunities with Massive Black Hole Binaries



Alberto Sesana
(Università di Milano Bicocca)



OUTLINE

Dynamics of MBH binary (MBHB) formation and dynamics

emission from MBHBs: gravitational waves (GWs) and electromagnetic (EM) radiation

Multimessenger astronomy with LISA and Athena (and LSST/Rubin)

Multimessenger astronomy with pulsar timing arrays (PTAs)

Observational facts

1- In all the cases where the inner core of a galaxy has been resolved (i.e. In nearby galaxies), a massive compact object (which I'll call Massive Black Hole, MBH for convenience) has been found in the centre.

2- MBHs must be the central engines of Quasars: the only viable model to explain this cosmological objects is by means of gas accretion onto a MBH.


3- Quasars have been discovered at $z \sim 7$, their inferred masses are $\sim 10^9$ solar masses!

THERE WERE 10^9 SOLAR MASS BHs
WHEN THE UNIVERSE WAS < 1 Gyr OLD!!!


MBH formation and evolution have profound consequences for GW astronomy



Structure formation in a nutshell




+




(From de Lucia et al. 2006)

(Ferrarese & Merritt 2000, Gebhardt et al. 2000)




(Menou et al 2001, Volonteri et al. 2003)

Structure formation in a nutshell




+



(From de Lucia et al. 2006)

(Ferrarese & Merritt 2000, Gebhardt et al. 2000)





- *Where and when do the first MBH seeds form?
- *How do they grow along the cosmic history?
- *What is their role in galaxy evolution?
- *What is their merger rate?
- *How do they pair together and dynamically evolve?

(Menou et al 2001, Volonteri et al. 2003)



Mergers







MBHB dynamics (BBR 1980)




MBHB dynamics (BBR 1980)









Milosavljevic & Merritt 2001





Colpi & Dotti 2009

MBHB dynamics (BBR 1980)





MBHB dynamics (BBR 1980)




But do we see them?

10 kpc: double quasars
(Komossa 2003)






1 kpc: double peaked NL
(Comerford 2013)




10 pc: double radio cores
(Rodriguez 2006)

1 pc: -shifted BL (Tsalmatzsa 2011)
-accelerating BL (Eracleous 2012)




0.01 pc: periodicity (Graham 2015)



0.0 pc: -X-shaped sources (Capetti 2001)
-displaced AGNs (Civano 2009)

characteristic amplitude




The Laser Interferometer Space Antenna

(LISA Consortium 2017)

Sensitive in the mHz frequency range where MBH binary evolution is fast (chirp)


Observes the full inspiral/merger/ringdown



Nicolas Douillet - ARTEMIS

3 satellites trailing the Earth connected through laser links


Proposed baseline:
2.5M km armlength
6 laser links
4 yr lifetime (10 yr goal)



The LISA Consortium

- Now a thriving community: 1300+ among full and associate members
- Several working groups connecting to the community: astrophysics, fundamental physics, cosmology, waveforms
- Several working packages defining deliverables
- 2 consortium meetings/yr, LISA symposium every 2 years, dedicated WG meetings every year

<https://www.lisamission.org/>



LISA Consortium User Guide

User guide ▾ Groups ▾ Getting help ▾ Contributing

Search Previous Next

LISA Consortium User Guide

This User Guide goal is to gather all the information related to the LISA Consortium tools. Users are more than welcome to contribute to its improvement. To do so, see the [HowToContribute](#) page.

- LISA Consortium User Guide
- Key information
- Collaborative tools
- Development tools and guidelines
- Sharing data tools
- Computing resources

LISA Consortium User Guide

User guide ▾ Groups ▾ Getting help ▾ Contributing

Search Previous

Key information

- LISA Consortium website
- Sign-up for the LISA Consortium
- Organisation
- LISA websites
- Key documents
- Next meetings (need to be logged to the wiki - see [LISA wiki](#))
- Acronyms
- Publication and Presentation Committee
- Inclusion and Diversity Committee
- Positions related to LISA

Collaborative tools

- LISA wiki
- LISA Document Management System (DMS) - Atrium
- Mailing lists
- Messaging on slack channels
- Audio / Video teleconferences

Development tools and guidelines

Mailing lists

- Management
- Full Member Groups
 - LISA Instrument Group
 - LISA Data Processing Group
 - LISA Science Group
 - Simulation Working Groups
- Associate and Full Members Groups
 - LISA Data Challenge Working Groups
 - Astrophysics Working Groups
 - Cosmology Working Groups
 - Fundamental Physics Working Groups
 - Waveform Working Groups
 - Advocacy and Outreach Working Groups

Mailing lists

- Consortium : consortium@lisamission.org


Management

- Consortium Lead : consortiumlead@lisamission.org
- Exec Board : exec_board@lisamission.org
- Board Member : board@lisamission.org
- Coordinator : coord@lisamission.org
- Coordination Group : coordination@lisamission.org
- Publication Committee : pubcom@lisamission.org
- Publication Committee Chairs : pubcom-chairs@lisamission.org

Full Member Groups

LISA Instrument Group


- LISA Instrument Group : lig@lisamission.org
- LIG Core : lig-core@lisamission.org
- LIG Performance Modelling WG : lig-pmwg@lisamission.org
- LIG-OB : lig-ob@lisamission.org
- LIG-PMS : lig-pms@lisamission.org
- LIG-GRS : lig-grs@lisamission.org
- LIG-OMS : lig-oms@lisamission.org
- LIG-Chairs : lig-chairs@lisamission.org
- LIG SLWG Chairs : lig-slwg-chairs@lisamission.org
- LIG Performance Modelling WG Chairs : lig-pmwg-chairs@lisamission.org







Associated electromagnetic signatures

In the standard circumbinary disk scenario, the binary carves a cavity: no EM signal (Phinney & Milosavljevic 2005).

However, all simulations (hydro, MHD) showed significant mass inflow (Cuadra et al. 2009, Shi et al 2011, Farris et al 2014, Tang et al. 2018...)





Simulations in hot gaseous clouds. Significant flare associated to merger (Bode et al. 2010, 2012, Farris et al 2012)




Simulations in disk-like geometry. Variability, but much weaker and unclear signatures (Bode et al. 2012, Gold et al. 2014)

Full GR force free electrodynamics (Palenzuela et al. 2010, 2012)




L_X




D'Ascoli et al. 2018

merger

post-merger



Disc re-brightening? Jet?




$t = -\infty$

$t = -20 \text{ h}$



$t = 0$

h_{gw}



(Palenzuela+ 2010, Gold+ 2014, Farris+ 2014, Tang+ 2017, 2018, D'Ascoli+ 2018, ...)

Opportunities for LISA-Athena (LSST/Rubin) synergies





Athena Wide Field Imager (WFI) (Rau+ 2015)

(Rau+ 2015)

Parameter	Characteristic
Energy Range	0.1-15 keV
Field of View	ca. 40' x 40' (baseline)

-X-ray telescope

-L2 ESA mission (~2030)



LSST : Vera Rubin observatory (Abell+ 2009)




-2022+


-Optical telescope

-9.6 square degree FoV


-m~24 within 30s pointings in several different filters

→ HOW CAN LISA AND ATHENA WORK TOGETHER?






Athena field of view





(Mangiagli+ 2020, Piro+ in prep.)

- Athena pre-pointing only possible for very low z sources
- LSST/Rubin more suitable for tracking inspiral periodicity (but optical)



1/5 of the
observable volume
of the universe.


Universe was
2Gyr old.




(McGee+ 2020)

Why multimessenger?


- Cosmology and cosmography at high z
- Study of accretion on MBHs with known mass and spins
- Study of the interplay between MBHs and gas (torques, disk structure, disk models)
- Host galaxy, Jet launches, Quasar birth ...



Courtesy of N. Tamanini



characteristic amplitude



Pulsar timing

Pulsars are neutron stars seen through their regular radio pulses

Pulsar timing is the art of measuring the time of arrival (ToA) of each pulse and then subtracting off the expected time of arrival given by a theoretical model for the system


1-Observe a pulsar and measure the ToAs

2-Find the model which best fits the ToAs

3-Compute the timing residual R


$$R = \text{ToA} - \text{ToA}_m$$

If the timing solution is perfect (and observations noiseless), then $R=0$. R contains all uncertainties related to the signal propagation and detection, plus the effect of unmodelled physics, like (possibly) *gravitational waves*









Earth




Pulsar




Incoherent superposition of sinusoids




$$h_c(f) = A \left(\frac{f}{\text{yr}^{-1}} \right)^{-2/3}$$



Simulated signal




Actual data




(Courtesy of PPTA)

Resolvable sources (AS et al 2009)



- *It is not smooth
- *It is not Gaussian
- *Single sources might pop-up
- *The distribution of the brightest sources might well be anisotropic

Finding the right galaxy





An individual PTA source must be massive and/or nearby → Only several tens of credible candidates (Goldstein et al 2019)

In general, PTA cannot break the distance-mass degeneracy ($A \sim M^{5/3}/D$)


$$A = 4 \frac{(G \mathcal{M}_z)^{5/3} (\pi f)^{2/3}}{D_1}$$

Sky localization is tens of deg^2 so tens of thousands of potential host galaxies



Associated electromagnetic signatures PTA


MBH binary + circumbinary disk



(Roedig et al. 2011, AS et al. 2012,
Tanaka et al. 2012, Burke-Spolaor 2013)

Associated electromagnetic signatures PTA

MBH binary + circumbinary disk



(Roedig et al. 2011, AS et al. 2012,

Tanaka et al. 2012, Burke-Spolaor 2013,

Farris+, D'Orazio+, Haiman+, Tang+,...)

A variety of possibilities:

Optical/IR dominated by the outer disk:
Steady/modulated?


UV generated by inner streams/minidisk:
periodic variability?

X rays variable from periodic shocks or intermittent corona?

Variable broad emission line in response to the varying ionizing continuum?


Double fluorescence lines?

Example: variability (AS+ 2012)



Streams feed the inner minidisk extremely intermittent mass inflow.

Applying this model to a typical MBH binary population we get ~ 100 sources at the eRosita flux limit



The future



MeerKAT, South Africa (2017)

The future



FAST, China (2017)

The future



Square Kilometre Array (SKA, 2021+)

Doggybag

MBHBs:

- are expected to form in the aftermath of galaxy mergers
- their dynamics is still a matter of active research, but binaries should form and coalesce within a Hubble time (reference figure: 10/yr)
- are the loudest GW sources in the Universe
- are expected to have an extravaganza of EM counterparts (but signatures?)

Joint GW-EM observations provide a number of benefits:

- Accretion physics
- Cosmography

LISA + Athena and/or LSST/Rubin might observe up to tens MBHBs in both GW and EM

PTA sources are massive and nearby, they might be 'easily' identified in the EM window

