

DT Invariants and Holomorphic Curves

Pierrick Bousseau

University of Georgia

Wall-Crossing Structures, Analyticity and Resurgence
IHES, 9 June 2023

Relation between two topics:

- Donaldson–Thomas (DT) invariants of non-compact Calabi–Yau 3-folds:
 - ▶ Counts of stable coherent sheaves (or complexes of coherent sheaves) on X or special Lagrangian submanifolds of its mirror Y .
 - ▶ Counts of BPS states for the $\mathcal{N} = 2$ 4d field theory T obtained by considering IIA or IIB string theory on $X \times \mathbb{R}^4$.
- Holomorphic curves in a hyperkähler manifold \mathcal{M} .
 - ▶ \mathcal{M} : Coulomb branch of T on $S^1 \times \mathbb{R}^3$, Seiberg–Witten integrable system.

- General expected picture [Kontsevich-Soibelman 1303.3253]
- A concrete example [B 1909.02985-1909.02992, B-Descombes-Le Floch-Pioline 2210.10712]:
 - ▶ DT invariants for coherent sheaves on local \mathbb{P}^2 : $X = K_{\mathbb{P}^2} = \mathcal{O}_{\mathbb{P}^2}(-3)$, non-compact Calabi-Yau 3-fold.
 - ▶ holomorphic curves in \mathcal{M} , (\mathcal{M}, I) : elliptic fibration, $(\mathcal{M}, J) = \mathbb{P}^2 \setminus E$, ALH^* metric.
 - ▶ Related to the expected resurgence properties of the solutions and quantum periods of a difference equation (quantum mirror curve).
- An heuristic/physics derivation of the general correspondence [B 2210.17001]
 - ▶ Holomorphic Floer theory for \mathcal{M} .

- DT invariants:

$$\Omega_{\gamma}(u) \in \mathbb{Z}$$

counts of geometric objects on a Calabi-Yau 3-fold X , with given topology class $\gamma \in \mathbb{Z}^n$ and satisfying a (Bridgeland) stability condition u .

- Examples:

- ▶ Stable holomorphic vector bundles of Chern character γ for a Kähler parameter u .
- ▶ Special Lagrangian submanifolds of class γ for a complex parameter u .

- $\mathcal{N} = 2$ supersymmetric 4d field theories
 - ▶ B : Coulomb branch of vacua of the 4d theory, $B \simeq \mathbb{C}^r$.
 - ▶ In a generic vacuum $u \in B \setminus \Delta$, abelian gauge theory $U(1)^r$
 - ▶ Supersymmetry: charge γ , central charge $Z_\gamma(u) \in \mathbb{C}$, BPS bound

$$|M| \geq |Z_\gamma(u)|$$

- ▶ Space of BPS states, saturating the BPS bound: $H_\gamma(u)$
- ▶ BPS index

$$\Omega_\gamma(u) = \text{Tr}_{H_\gamma(u)}(-1)^F$$


- Geometric constructions from string theory: IIA or IIB string on Calabi-Yau 3-fold X
- Expectations:
 - ▶ the universal cover of $B \setminus \Delta$ naturally maps to the space of Bridgeland stability conditions.
 - ▶ DT invariants = BPS indices: stability $u \in B \setminus \Delta$
- From now on: consider $\mathcal{N} = 2$ 4d field theories without gravity.
 - ▶ Geometrically: non-compact Calabi-Yau 3-folds.

Wall-crossing

- $\Omega_\gamma(u)$: constant function of u away from codimension one loci in B , called walls, across which $\Omega_\gamma(u)$ jumps discontinuously.
- Jumps controlled by a universal wall-crossing formula [Kontsevich-Soibelman]:

$$\{\Omega_\gamma(u^-)\}_\gamma \rightarrow \{\Omega_\gamma(u^+)\}_\gamma.$$

- Example: $\mathcal{N} = 2$ $SU(2)$ gauge theory



Seiberg-Witten integrable system


- \mathcal{M} : Coulomb branch of the theory on $\mathbb{R}^3 \times S^1$, hyperkähler manifold of complex dimension $2r$, complex integrable system:

$$\pi: \mathcal{M} \longrightarrow B$$

- Twistor sphere of complex structures I, J, K
 - ▶ π I -holomorphic: in complex structure I , generic fibers of π are abelian varieties of dimension r .
 - ▶ for every $\theta \in \mathbb{R}/2\pi\mathbb{Z}$, generic fibers of π are special Lagrangians in complex structure $J_\theta = (\cos \theta)J + (\sin \theta)K$.
 - ▶ Low energy: 3d $\mathcal{N} = 4$ sigma model with target \mathcal{M}
- $u \in B \setminus \Delta$, $\gamma \in \pi_2(\mathcal{M}, \pi^{-1}(u)) \rightarrow H_1(\pi^{-1}(u), \mathbb{Z}) = \mathbb{Z}^{2r}$,

$$Z_\gamma(u) = \int_\gamma \Omega_I$$

Seiberg-Witten integrable system




- Class S on C : $\pi: \mathcal{M} \rightarrow B$ is (essentially) the Hitchin integrable system for C .

- For every point $u \in B \setminus \Delta$, and class $\gamma \in \pi_2(\mathcal{M}, \pi^{-1}(u))$,

$$\Omega_\gamma(u) = N_\gamma(u).$$

- ▶ $\Omega_\gamma(u)$: DT/BPS invariants counting u -stable objects of class γ .
- ▶ $N_\gamma(u)$: count of J_θ -holomorphic disks in \mathcal{M} with boundary on the fiber $\pi^{-1}(u)$ and of class γ , where $\theta = \text{Arg} Z_\gamma(u)$.
- Evidence:
 - ▶ BPS spectrum $\{\Omega_\gamma(u)\} \rightarrow$ hyperkähler geometry of \mathcal{M} [Gaiotto-Moore-Neitzke]
 - ▶ J_θ -holomorphic disks: instantons/quantum corrections to construct the mirror of $(\mathcal{M}, \omega_\theta)$ [Fukaya, Kontsevich-Soibelman, Gross-Siebert]
 - ▶ Same wall-crossing formula [Kontsevich-Soibelman]
 - ▶ Tropical curves in B from holomorphic disks and attractor trees from DT invariants [Kontsevich-Soibelman]

Expectation



Problems:

- The embedding of B in the space of Bridgeland stability conditions is not known in general.
- Defining counts of holomorphic disks is difficult in general.


Examples:

- Log Gromov–Witten invariants: algebro-geometric version of holomorphic disks used by Gross–Siebert in their mirror symmetry construction.
- DT invariants of quivers with potential versus log Gromov–Witten invariants of toric and cluster varieties [B–Argüz 2302.02068].
- This talk:
 - ▶ DT invariants counting coherent sheaves on local \mathbb{P}^2
 - ▶ One of the few examples where the embedding in the space of Bridgeland stability conditions is known.


- $X = K_{\mathbb{P}^2} = \mathcal{O}_{\mathbb{P}^2}(-3)$ non-compact Calabi-Yau 3-fold
 - ▶ Zero section $\iota: \mathbb{P}^2 \hookrightarrow X$
- $D_{\mathbb{P}^2}(X)$: bounded derived category of sheaves on X set-theoretically supported on \mathbb{P}^2
 - ▶ $\iota_*: D^b \text{Coh}(\mathbb{P}^2) \rightarrow D_{\mathbb{P}^2}(X)$
 - ▶ $\mathcal{O}(n) := \iota_* \mathcal{O}_{\mathbb{P}^2}(n)$ (D4-branes with n units of D2-charges)
- IIA string theory on X : $\mathcal{N} = 2$ 4d theory.
 - ▶ Seiberg-Witten geometry $\pi: \mathcal{M} \rightarrow B$?
 - ▶ Mirror symmetry: $B \setminus \Delta = \mathbb{H}/\Gamma_1(3)$, modular curve. \mathcal{M} : universal family of elliptic curves.

$$\mathcal{M} \rightarrow B$$

A fundamental domain F_C of $\Gamma_1(3)$ acting on \mathbb{H} :




The modular curve $B \setminus \Delta = \mathbb{H}/\Gamma_1(3)$:



$$\mathcal{M}' \rightarrow B'$$

- Work on the 3:1 cover B' of B resolving the orbifold point.
- $\mathcal{M}' \rightarrow B'$: elliptic fibration with 3 singular fibers.



Map to the space of stability conditions

- $Stab(D_{\mathbb{P}^2}(X))$: space of Bridgeland stability conditions on $D_{\mathbb{P}^2}(X)$, complex manifold of dimension 3
- Bayer-Macri (2009):

$$\widetilde{B \setminus \Delta} = \mathbb{H} \rightarrow Stab(D_{\mathbb{P}^2}(X))$$

$$\tau \mapsto (\mathcal{A}(\tau), Z(\tau))$$

- Central charge, additive map:

$$Z(\tau) : \Gamma = K_0(D_{\mathbb{P}^2}(X)) = \mathbb{Z}^3 \rightarrow \mathbb{C}$$

$$\gamma \mapsto Z_\gamma(\tau)$$

- To summarize:

$$\widetilde{B \setminus \Delta} = \mathbb{H} \rightarrow \text{Stab}(D_{\mathbb{P}^2}(X))$$
$$\tau \mapsto (\mathcal{A}(\tau), Z(\tau))$$

- We can then do DT theory.

- ▶ Moduli spaces

$$M(\gamma, \tau) = \{\tau\text{-semistable objects in } \mathcal{A}(\tau) \text{ of class } \gamma\}$$

- ▶ DT/BPS invariants:

$$\Omega(\gamma, \tau) \in \mathbb{Z}$$

- ▶ Wall-crossing as a function of $\tau \in \mathbb{H}$.

- Mirror curve:

$$H(p, x) = -1 + e^x + e^p + z(\tau) e^{-x-p} = 0,$$

- Quantum mirror curve, difference equation:

$$(-1 + e^x)\Psi(x) + \Psi(x + \hbar) + z(\tau) e^{\frac{\hbar}{2}} e^{-x} \Psi(x - \hbar) = 0.$$

- WKB solution: $\Psi(x) = \exp(\frac{1}{\hbar} \int^x \lambda(\hbar))$, where $\lambda(\hbar)$ is a 1-form with formal dependence on \hbar : $\lambda(\hbar) = \sum_{j \geq 0} \lambda_j \hbar^j$.
- Formal “Quantum periods”

$$a(\hbar) = \int_{\gamma} \lambda(\hbar), \quad a_D(\hbar) = \int_{\gamma_D} \lambda(\hbar).$$


- Expectation: resurgence of the quantum periods is controlled by the BPS spectrum $\{\Omega(\gamma, \tau)\}_{\gamma}$. [Gu-Marino 2211.03871]

Scattering diagrams

- Goal: study of the DT/BPS invariants using flow trees organized in “scattering diagrams” in $\widetilde{B \setminus \Delta} = \mathbb{H}$
- Pick a phase $\theta \in \mathbb{R}/2\pi\mathbb{Z}$
 - For every $\gamma \in \Gamma$, consider the 1-dimensional locus, “rays”:

$$\mathcal{R}_\gamma^+(\theta) := \{\tau \in \mathbb{H} \mid \text{Arg}(Z_\gamma(\tau)) = \theta, \Omega(\gamma, \tau) \neq 0\} \subset \mathbb{H}$$

- Orient rays such that $|Z_\gamma(\tau)|$ increases.
- Decorate the rays by generating functions of DT invariants, get a scattering diagram \mathfrak{D}_θ




Theorem (B, Descombes, Le Floch, Pioline, 2022)

For every $\theta \in \mathbb{R}/2\pi\mathbb{Z}$, the scattering diagram \mathcal{D}_θ can be uniquely reconstructed from:

- *Explicit initial rays coming from the conifold points.*
- *Scatterings imposed by the consistency condition.*
- Algorithmic reconstruction of the full BPS spectrum (except pure D0) at any point of the physical space of stability conditions.


Initial rays

At the conifold point $\tau_O = 0$, $Z_O(\tau_O) = 0$. Infinitely many initial rays corresponding to the objects $\mathcal{O}[k]$, $k \in \mathbb{Z}$.




General conifold point: apply $\Gamma_1(3)$, spherical object E becoming massless, infinitely many initial rays corresponding to the objects $E[k]$, $k \in \mathbb{Z}$.

The scattering diagram $\mathfrak{D}_{\frac{\pi}{2}}$



The scattering diagram \mathfrak{D}_0




- Expectation: for every $\theta \in \mathbb{R}/2\pi\mathbb{Z}$, the scattering diagram \mathfrak{D}_θ should describe J_θ -holomorphic disks in \mathcal{M}' .
- Problem: how to describe \mathcal{M}' as a complex manifold for the complex structure J_θ ?
 - ▶ We only know that (\mathcal{M}', I) is an elliptic fibration over B .
- [Collins-Jacob-Lin]:
 - ▶ $(\mathcal{M}', J_{\frac{\pi}{2}}) = \mathbb{P}^2 \setminus E$, where $E \subset \mathbb{P}^2$ is a smooth cubic. Affine algebraic variety.
 - ▶ $(\mathcal{M}', J_0) \simeq (\mathcal{M}', I)$, elliptic fibration. Twin torus fibrations.
- In both cases, use algebro-geometric definition of counts of holomorphic disks as log Gromov–Witten invariants.

Holomorphic disks?

Theorem (Gräfnitz, B.)

The scattering diagram $\mathfrak{D}_{\frac{\pi}{2}}$ describes log curves in $(\mathcal{M}', J_{\frac{\pi}{2}}) = \mathbb{P}^2 \setminus E$.



Corollary (B.)

Correspondence between DT invariants of $K_{\mathbb{P}^2}$ of phase $\frac{\pi}{2}$ and counts of log curves in $\mathbb{P}^2 \setminus E$


Applications:

- Proof of Takahashi's conjecture on Gromov–Witten invariants of (\mathbb{P}^2, E) [B].
- Proof of quasimodularity of generating series of DT invariants [B-Fan-Guo-Wu].

The scattering diagram \mathfrak{D}_0

Theorem (Gross-Hacking-Keel)

The scattering diagram \mathfrak{D}_0 describes log curves in (\mathcal{M}', J_0) .



Corollary (B.)

Correspondence between DT invariants of $K_{\mathbb{P}^2}$ of phase 0, DT invariants of the quiver (Q, W) , and counts of log curves in the elliptic fibration (\mathcal{M}', J_0) .

Why are the counts of BPS states of a $\mathcal{N} = 2$ 4d theory given by counts of holomorphic curves in the Seiberg–Witten geometry $\pi : \mathcal{M} \rightarrow B$?

- Mirror symmetry and hyperkähler rotation for $X = K_{\mathbb{P}^2}$.
- In general?
 - ▶ Stronger conjecture formulated using holomorphic Floer theory.
 - ▶ Physics derivation.

- How to go from coherent sheaves on $X = K_{\mathbb{P}^2}$ to J_θ -holomorphic curves in the Coulomb branch $\pi: \mathcal{M} \rightarrow B$?
- Claim: the mirror Y of X is the non-compact Calabi-Yau 3-fold $Y: uv = \pi - t$.
 - ▶ Hyperkähler rotation: J_θ -holomorphic disks in $\mathcal{M} \rightarrow$ special Lagrangian disks in (\mathcal{M}, I) of phase θ .
 - ▶ Suspension \rightarrow closed special Lagrangians in Y .
 - ▶ Mirror symmetry \rightarrow stable coherent sheaves on X .
- Physics: IIA on $X \leftrightarrow$ IIB on $Y \leftrightarrow$ IIA on \mathcal{M} and NS5 on $\pi^{-1}(u) \leftrightarrow$ M on \mathcal{M} and M5 on $\pi^{-1}(u) \leftrightarrow$ IIB on B , D3 on u (string junctions on D3-brane probe)

[Kontsevich-Soibelman], [Doan-Rezchikov], [B]


- $(\mathcal{M}, I, \Omega_I)$: holomorphic symplectic manifold.
 - ▶ Hyperkähler structure I, J, K , $J_\theta := (\cos \theta)J + (\sin \theta)K$.
 - ▶ $L_1, L_2 \subset \mathcal{M}$: I -holomorphic Lagrangian, $\Omega_I|_{L_1} = \Omega_I|_{L_2} = 0$.
- P : space of paths between L_1 and L_2 , $W := \int_p d^{-1}\Omega_I$ (multivalued!)
 - ▶ Critical points: intersection points $L_1 \cap L_2$.
 - ▶ Gradient flow lines: J_θ holomorphic curves, $u : \mathbb{R}^2 \rightarrow \mathcal{M}$.
 - ▶ ζ -instantons, $u : \mathbb{R}^3 \rightarrow \mathcal{M}$, solutions to Fueter equation

$$\partial_\tau u + I\partial_s u + J_\theta \partial_t u = 0.$$


- LG model for (P, W) :
 - ▶ $p, q \in L_1 \cap L_2 \rightarrow$ vector space H_{pq} of 2d BPS states of (P, W)
 - ▶ $L_1, L_2 \rightarrow$ category $\text{Brane}(P, W)$
 - ▶ $\mathcal{M} \rightarrow$ 2-category of I -holomorphic Lagrangians (A-model versus Rozansky-Witten B-model).

Holomorphic Floer theory and DT invariants

- Back to a $\mathcal{N} = 2$ 4d field theory.
 - How to recover the BPS spectrum $\{\Omega_\gamma(u)\}$ from holomorphic Floer theory? Correct holomorphic symplectic manifolds \mathcal{M} and holomorphic Lagrangians L_1, L_2 ?
 - ▶ \mathcal{M} : Seiberg-Witten integrable system
 - ▶ $L_1 = \pi^{-1}(u)$: fiber of $\pi : \mathcal{M} \rightarrow B$ over $u \in B$.
 - ▶ $L_2 = S$: natural section of π . Physical definition: boundary condition for the 3d sigma model of target \mathcal{M} defined by the cigar geometry
- [Nekrasov-Witten] Hitchin system example: Hitchin section.




Holomorphic Floer theory and DT invariants



- $L_1 \cap L_2 = \pi^{-1}(u) \cap S = \{p\}$
- But $\pi_1(P) \neq 0$ and W is multivalued.
- $\pi_1(P) = \pi_2(\mathcal{M}, \pi^{-1}(u))$: on \tilde{P} , critical points of W indexed by


$$\gamma \in \pi_2(\mathcal{M}, \pi^{-1}(u))$$




Conjecture (B)

Given a $\mathcal{N} = 2$ 4d field theory, the space of BPS states $H_\gamma(u)$ of class γ in the vacuum u is isomorphic to the vector space $H_{0\gamma}$ associated by holomorphic Floer theory for the Seiberg-Witten integrable system \mathcal{M} to the lifts 0 and γ of the intersection point between the fiber $\pi^{-1}(u)$ and the section S :

$$H_\gamma(u) \simeq H_{0\gamma}$$



Gradient flow lines are naturally J_θ -holomorphic disks with boundary on $\pi^{-1}(u)$ and so one recovers the previous expectation in the numerical limit.



Thank you for your attention !